

Estimates of First and Second Order Shape Derivatives in Nonsmooth Multidimensional Domains and Applications

Jimmy Lamboley*, Arian Novruzi†, Michel Pierre‡

February 24, 2016

Abstract

In this paper we investigate continuity properties of first and second order shape derivatives of functionals depending on second order elliptic PDE's around nonsmooth domains, essentially either Lipschitz or convex, or satisfying a uniform exterior ball condition. We prove rather sharp continuity results for these shape derivatives with respect to Sobolev norms of the boundary-traces of the displacements. With respect to previous results of this kind, the approach is quite different and is valid in any dimension $N \geq 2$. It is based on sharp regularity results for Poisson-type equations in such nonsmooth domains. We also enlarge the class of functionals and PDEs for which these estimates apply. Applications are given to qualitative properties of shape optimization problems under convexity constraints for the variable domains or their complement.

MSC2010: 49Q10, 46N10, 46E35, 35R35.

Keywords: Shape derivative, shape optimization, convexity constraint, energy functional, Sobolev estimates, optimality conditions, regularity.

1 Introduction

In this paper, we focus on regularity estimates for first and second order shape derivatives *around nonsmooth subsets of \mathbb{R}^N ($N \geq 2$)* for energy functionals involving classical elliptic partial differential equations (PDE). For instance, we address the following question: given a bounded Lipschitz or convex subset $\Omega_0 \subset \mathbb{R}^N$, we wonder what is the optimal regularity of the shape derivatives

$$\xi \rightarrow E'(\Omega_0)(\xi), \quad \xi \rightarrow E''(\Omega_0)(\xi, \xi),$$

where $E'(\Omega_0), E''(\Omega_0)$ respectively denote the first and second shape derivatives around Ω_0 of the shape functional $\Omega \mapsto E(\Omega) = \int_{\Omega} K(x, U_{\Omega}, \nabla U_{\Omega}) dx$, $K(x, \cdot, \cdot)$ a quadratic polynomial and $U_{\Omega} = U_{\Omega}(x)$ the solution of an elliptic PDE set in Ω (see Sections 2.1, 2.2 for the precise definitions).

This question, interesting for itself, is in particular motivated by the qualitative analysis of shape optimization problems of the form

$$\min\{J(\Omega), \Omega \subset \mathbb{R}^N \text{ convex}, \Omega \in \mathcal{O}_{ad}\}, \tag{1}$$

where \mathcal{O}_{ad} is a set of admissible subsets of \mathbb{R}^N and $J : \mathcal{O}_{ad} \rightarrow \mathbb{R}$ is a shape functional which itself involves shape functionals $\Omega \mapsto E(\Omega)$ of the above type.

*CEREMADE, Université Paris-Dauphine, Place du Maréchal de Lattre de Tassigny, 75775 Paris, France

†University of Ottawa, Department of Mathematics and Statistics, 585 King Edward, Ottawa, ON, K1N 6N5, Canada

‡ENS Rennes, IRMAR, UEB, av Robert Schuman, 35170 Bruz, France

The following 2-dimensional example was considered in [20] among other cases:

$$J(\Omega) = R(E(\Omega), |\Omega|) - P(\Omega), \quad \mathcal{O}_{ad} = \{\Omega \subset \mathbb{R}^2, \text{ open and } \partial\Omega \subset \{x, a \leq |x| \leq b\}\}, \quad (2)$$

where $R : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a smooth function, $E(\Omega)$ is a shape functional related to a PDE, $|\Omega|$ is the Lebesgue measure of Ω , $P(\Omega)$ its perimeter and $(a, b) \in [0, \infty]^2$. It was proved (see [20, Theorem 2.9, Theorem 2.12 and Corollary 2.13] and also Section 4 in the present paper) that if the second order shape derivative $\xi \mapsto E''(\Omega_0)(\xi, \xi)$ around any bounded convex subset Ω_0 is *continuous with respect to a norm strictly weaker than $H^1(\partial\Omega_0)$* , then optimal shapes of (1) are *polygonal* in $\{x, a < |x| < b\}$. In [20], the authors prove that such a continuity holds in the two specific examples: when the functional $E(\Omega)$ is the Dirichlet energy of Ω -that is $K = K(x, U, q) = \|q\|^2 - 2f(x)$ and U_Ω is solution of the associated Dirichlet problem, or when $E(\Omega)$ is the first eigenvalue of the Dirichlet-Laplacian on Ω . More precisely, it is proved in [20] that the second order shape derivative of $E(\cdot)$ is, in these two examples and around any open convex domain Ω_0 , continuous for the $H^{1/2}(\partial\Omega_0) \cap L^\infty(\partial\Omega_0)$ topology (and therefore continuous for the $H^{1/2+\varepsilon}(\partial\Omega_0)$ -topology for any $\varepsilon > 0$). The proof of this continuity strongly relies on the 2-dimensional environment. As explained below, we prove in this paper that *even the full $H^{1/2}(\partial\Omega_0)$ -continuity holds in this case and even in any dimension* (see iii) in Corollary 2.6). Note that this continuity is optimal since, for regular convex domains Ω_0 , if for instance $f = 0$ on $\partial\Omega_0$, then $E''(\Omega_0)$ satisfies

$$E''(\Omega_0)(\xi, \xi) \geq C \|\xi\|_{H^{1/2}(\partial\Omega_0)}^2, \quad (3)$$

for any displacement ξ which is orthogonal to $\partial\Omega_0$. This estimate (3) may be obtained by using (5.101) and Section 5.9.6. in [12].

In Problem (1), only convex domains are involved. However, it is also interesting to consider shape optimization problems where the PDE is set in an "exterior domain" like

$$\min\{J(\mathbb{R}^N \setminus \overline{\Omega}), \Omega \subset \mathbb{R}^N \text{ convex}, \Omega \in \mathcal{O}_{ad}\}, \quad (4)$$

where \mathcal{O}_{ad} is as before, but now J involves $E(\Omega) = \int_{\mathbb{R}^N \setminus \overline{\Omega}} K(U_\Omega, \nabla U_\Omega)$ where U_Ω is solution of an elliptic PDE set on the *exterior domain* $\mathbb{R}^N \setminus \overline{\Omega}$, which is here the complement of a convex set. It is well-known that solutions of such PDE's are not so regular as in convex domains. These exterior domains are nevertheless Lipschitz domains and this is one main reason why it is interesting to look at the case of Lipschitz domains even for shape optimization problems involving convex domains like (4).

In this paper, we use a different strategy to estimate shape derivatives, and we generalize and improve the above-mentioned shape derivative estimates in the following directions.

- A most important generalization concerns the dimension of shapes. In [20], the result and the strategy were restricted to **planar shapes**. Thanks to our new strategy, we are able to provide estimates of first and second order shape derivatives **in any dimension**. This is an important breakthrough for the study of problems like (1) in dimension 3 or higher.
- We generalize the class of PDE energy-functionals in several ways (see Section 2.1 for precise definitions): the underlying elliptic operator is now a general linear elliptic operator with variable coefficients, the energy is any quadratic integral functional of the state function and its gradient (and in particular does not need to be the energy associated to the PDE defining the state function) and more importantly, we consider both interior and exterior PDEs. This last point motivates the next item.
- We generalize the classes of shapes we consider. Indeed we do not only consider the class of convex domains, but we investigate two wider classes, namely the class of Lipschitz domains and the class

of Lipschitz domains satisfying a uniform exterior ball condition (we refer to these domains as *semi-convex* domains, see Definition 2.1). Even if we are interested in applications about convex domains, as we pointed out in the previous point, we are interested in PDEs defined in the exterior of some convex domain. Whence the consideration of the above kinds of boundary regularity. Estimates for semi-convex domains are the same as those for convex domains. They are weaker for Lipschitz domains, but probably rather sharp and they are strong enough to be used for our applications (see Section 4) which are interesting for themselves.

- Even in the particular case where E is exactly the Dirichlet energy and Ω is a 2-dimensional convex domain, we improve the result of [20] in obtaining that $E''(\Omega)$ is continuous in the $H^{1/2}(\partial\Omega)$ -norm, instead of $H^{1/2}(\partial\Omega) \cap L^\infty(\partial\Omega)$. As noticed above, this result is sharp since one cannot expect continuity in an H^s -norm for $s < 1/2$ (even if Ω is smooth). This specific result is actually valid as soon as the energy is the one associated to the PDE defining the state function, and is valid in any dimension and for any semi-convex domain.

The method for proving the shape derivative estimates is new. In order to study the derivative at a set Ω_0 , we first prove adequate estimates of the “material derivative” \hat{U}'_θ , where $\theta : \mathbb{R}^N \rightarrow \mathbb{R}^N$ is a smooth vector field, $'$ denotes the derivative with respect to θ , $\hat{U}_\theta = U_\theta \circ (I + \theta)$, and U_θ is the state solution of the related PDE in the domain $\Omega_\theta = (I + \theta)(\Omega_0)$. For our purpose, it appears that it is much more efficient than using the usual shape derivative U'_θ . We use the well-known property (see for example [12]) that without any regularity of Ω , the map $\theta \mapsto \hat{U}_\theta$ is in general differentiable (even C^∞ if the involved coefficients are smooth) when seen as a $H_0^1(\Omega_0)$ -valued function (whereas $\theta \mapsto U'_\theta$ is differentiable only when seen as an L^2 -valued function). We show how the regularity of the shape derivatives essentially depends on the regularity of the state function U_0 . Thus we shall use some sharp regularity results for the solution of a linear second order PDE in a Lipschitz or semi-convex domain, in particular $W^{1,p}$ -regularity results when the data are in $W^{-1,p}$ (see Propositions 3.3 and 3.5).

Note that we insisted in this introduction on the second order shape derivative, but we also obtain estimates of the first order shape derivative which seem to be new as well, in this non-smooth setting.

As an application of these estimates, using the strategy of [19, 20], we prove that any solution Ω_0 of (1), (4) is polygonal, if $N = 2$ and if E is in one of two classes described above (where E depends on the solution of a PDE in the interior or the exterior of a convex domain, see more precisely Section 2.1). Also, in higher dimension, we use our estimates to analyze solutions of the N -dimensional version of Problem (2). We obtain very strong qualitative properties of optimal shapes, namely that the space of deformations which leave it convex is actually *of finite dimension* (see Theorem 4.8). As an easy consequence, we obtain that any optimal shape has zero Gauss curvature on any open set where the boundary is smooth. Actually, this “finite dimension” property does contain quite more geometrical information.

This paper is structured as follows. In Section 2 we introduce the notations, the classes of PDE’s under consideration and state our main results. In Section 3 we prove the estimates of first and second order shape derivatives. In the last section, we apply these estimates for the analysis of optimal planar convex domains, which happen to be polygons and we conclude by analyzing the consequence of our estimates for higher dimensional optimal convex domains.

2 Main results

2.1 Class of energy functionals

We deal with energy functionals $E(\Omega)$ of two main forms.

- **Interior PDE:** $E(\Omega)$ depends on a PDE in the interior of Ω , namely

$$E(\Omega) = \int_{\Omega} K(x, U_{\Omega}, \nabla U_{\Omega}) dx, \quad (5)$$

where

$$K(x, U, q) = \frac{1}{2} \left(\sum_{i,j=1}^N \alpha_{ij}(x) q_i q_j + \alpha_{00}(x) U^2 \right) + \sum_{i=1}^N \beta_i(x) q_i + \gamma(x) U + \delta(x), \quad (6)$$

with smooth enough coefficients $\alpha, \beta, \gamma, \delta$, and U_{Ω} solution of

$$U \in H_0^1(\Omega), \quad LU := - \sum_{i,j=1}^N \partial_j (a_{ij}(x) \partial_i U) + \sum_{i=1}^N b_i \partial_i U + cU = f, \quad \text{in } \Omega. \quad (7)$$

Regularity of the coefficient will be made precise later, but they will satisfy the following throughout the paper:

$$\exists \lambda > 0, \quad \forall \xi \in \mathbb{R}^N, \quad \forall x \in \mathbb{R}^N, \quad \lambda |\xi|^2 \leq \sum_{i,j=1}^N a_{ij}(x) \xi_i \xi_j \leq \lambda^{-1} |\xi|^2, \quad (8)$$

$$\forall (i, j) \in \llbracket 1, n \rrbracket, \quad a_{ij} = a_{ji}, \quad \alpha_{ij} = \alpha_{ji}, \quad \text{and } c \geq 0. \quad (9)$$

Note that the condition $\alpha_{ij} = \alpha_{ji}$ is not restrictive as, in the case the matrix (α_{ij}) is not symmetric, we can consider $\bar{\alpha}_{ij} = \frac{1}{2}(\alpha_{ij} + \alpha_{ji})$, which is symmetric, and then in (6) take $\bar{\alpha}_{ij}$ instead of α_{ij} . Note also that no ellipticity condition is a priori required on (α_{ij}) . Actually, our strategy can handle much more general functional K , see Remark 3.15.

- **Exterior PDE:** For simplicity, we will restrict ourselves to the model example where

$$E(\Omega) = \int_{\Omega^e} |\nabla U_{\Omega}|^2, \quad (10)$$

where $U_{\Omega} \in H_0^{1,0}(\Omega^e)$ solves

$$-\Delta U = f \text{ in } \Omega^e := \mathbb{R}^N \setminus \bar{\Omega}. \quad (11)$$

and $H_0^{1,0}(\Omega^e)$ is defined (see [26]) as the set of functions of $H^{1,0}(\Omega^e)$ with zero trace on $\partial\Omega$ and

$$H^{1,0}(\Omega^e) = \left\{ U : \Omega^e \mapsto \mathbb{R} \text{ measurable, } \|U\|_{H^{1,0}(\Omega^e)}^2 := \int_{\Omega^e} \frac{U^2}{(1+|x|^2)[\ln(2+|x|^2)]^{\delta_{2,N}}} + |\nabla U|^2 < \infty \right\} \quad (12)$$

with $\delta_{2,N} = 1$ if $N = 2$ and $\delta_{2,N} = 0$ if $N \neq 2$.

2.2 Shape derivative estimates

In the whole paper, we consider

$$\Omega_0, \Omega \Subset B_R = B(0, R), \text{ connected, open, bounded,} \quad (13)$$

where $R > 0$ (and large). We set $\Theta := W_0^{1,\infty}(B_R, \mathbb{R}^N)$ the Banach space equipped with the usual norm

$$\forall \theta \in \Theta, \|\theta\|_{1,\infty} := \sup_{x \in B_R} \{\|\theta(x)\| + \|D\theta(x)\|\}.$$

Given a shape functional $E(\cdot) : \mathcal{O}_{ad} \rightarrow \mathbb{R}$ defined on a family \mathcal{O}_{ad} of admissible subsets of \mathbb{R}^N , we consider

$$\mathcal{E} : \Theta \rightarrow \mathbb{R}, \quad \mathcal{E}(\theta) := E((I + \theta)(\Omega_0)).$$

Then, E is said to be shape differentiable of order $m \in \mathbb{N}^*$ at Ω_0 (resp. of class C^m near Ω_0) if and only if \mathcal{E} is m times Fréchet-differentiable at $\theta = 0$ (resp. if and only if \mathcal{E} is m times continuously differentiable in a neighborhood of $\theta = 0$). When it is well-defined, $\mathcal{E}'(0)$ and $\mathcal{E}''(0)$ are respectively called the first and second order shape derivative of E at Ω_0 , and can also be denoted by $E'(\Omega_0)$ and $E''(\Omega_0)$. Their values at $\xi, \eta \in \Theta$ are denoted by $\mathcal{E}'(0)(\xi)$ or $E'(\Omega)(\xi)$, $\mathcal{E}''(0)(\xi, \eta)$ or $E''(\Omega)(\xi, \eta)$. They are linear and bilinear continuous forms on Θ respectively and it is well-known that $E'(\Omega_0), E''(\Omega_0)$ only depend on the trace on $\partial\Omega_0$ of the deformations ξ and η (see the proof of Theorem 2.3).

Definition 2.1 *A bounded open subset $\Omega \subset \mathbb{R}^N$ is said to be semi-convex if it is Lipschitz and satisfies a uniform exterior ball condition in the following sense: there exists $r > 0$ such that for any $x \in \partial\Omega$, there exists $y \in \mathbb{R}^N$ with $\overline{B(y, r)} \cap \Omega = \{x\}$.*

It is known that a domain is semi-convex if it is locally representable as the graph of a semi-convex function (see for example [23, Theorem 3.9]), where a function $f : C \rightarrow \mathbb{R}$ is said semi-convex on a convex subset C of \mathbb{R}^n if there exists $M \in \mathbb{R}$ such that $x \in C \mapsto f(x) + M\|x\|^2$ is convex. Recall that Ω is said to be Lipschitz if it is locally representable as the graph of a Lipschitz function.

Remark 2.2 It is easy to check that, if Ω is semi-convex then for R_1 small enough, there exist $(\delta_1(R_1), \sigma(R_1)) \in (0, \infty) \times (0, 1)$ with $\lim_{R_1 \rightarrow 0} \delta_1(R_1) = 0$ such that Ω is a $(\delta_1(R_1), \sigma(R_1), R_1)$ -quasi-convex domain in the sense of [14]. It follows that Ω is a (δ, σ, R) -quasi-convex domain in the sense of [14] for all $(\delta, \sigma, R) \in [\delta_1(R_1), \infty) \times (0, k(R)\sigma(R_1)] \times (0, R_1]$, with $k(R) > 0$, $k(R_1) = 1$, $\lim_{R \rightarrow 0} k(R) = 0$.

On the other hand, given a Lipschitz matrix (a_{ij}) (i.e. $a_{ij} \in W^{1,\infty}(\Omega) \forall i, j = 1, \dots, N$), then for all $R_2 \in (0, 1)$, there exists $\delta_2(R_2) \in (0, \infty)$ with $\lim_{R_2 \rightarrow 0} \delta_2(R_2) = 0$ such that A is a $(\delta_2(R_2), R_2)$ -vanishing matrix in the sense of [14]. It follows that A is (δ, R) -vanishing matrix in the sense of [14] for all $(\delta, R) \in [\delta_2(R_2), \infty) \times (0, R_2]$.

Therefore, we will be able to apply the $W^{1,p}$ -regularity results proved in Theorem 1.1 of [14] for the solutions to (7) in quasi-convex domains (see Proposition 3.5). Indeed, given $p \in (1, \infty)$, we first choose R_1 and R_2 small enough so that $\delta_1(R_1), \delta_2(R_2) \leq \delta(N, p)$ as defined in Theorem 1.1 of [14], and Ω is a $(\delta_1(R_1), \sigma(R_1), R_1)$ -quasi-convex and (a_{ij}) is $(\delta(R_2), R_2)$ -vanishing. It follows that if $R = \min\{R_1, R_2\}$ and $\sigma = k(R)\sigma(R_1)$ then Ω is a $(\delta(N, p), \sigma, R)$ -quasi-convex and (a_{ij}) is $(\delta(N, p), R)$ -vanishing.

Let us now present the main results of this paper.

Theorem 2.3 (interior) *Let Ω_0 be as in (13). For $\theta \in \Theta$, we denote by U_θ the solution of (7) in $\Omega_\theta = (I + \theta)(\Omega_0)$ (see Proposition 3.3) and $\mathcal{E}(\theta) = E(\Omega_\theta)$, where E is given by (5). The following holds.*

i) If $a_{ij}, b_i, c, f, \alpha_{ij}, \beta_i, \gamma, \delta \in W^{m,\infty}(B_R)$, $m \in \mathbb{N}^$, then $[\theta \mapsto \mathcal{E}(\theta)]$ is of class C^m near $\theta = 0 \in \Theta$.*

In the rest of this statement, we assume the previous hypotheses are satisfied for $m = 2$.

ii) Assume Ω_0 is Lipschitz and L is of the form (7). Then there exists $\bar{r}_1 = \bar{r}_1(\Omega_0)$ satisfying $\bar{r}_1 \in (1, 2)$ if $N = 2$ and $\bar{r}_1 \in (1, 3)$ if $N \geq 3$ such that, for all $r \in (\bar{r}_1, \infty)$, there exist $C_i = C_i(\Omega_0, L, f, K, r)$ with

$$|E'(\Omega_0)(\xi)| \leq C_1 \|\xi\|_{W^{1-1/r, r}(\partial\Omega_0)}, \quad (14)$$

$$|E''(\Omega_0)(\xi, \xi)| \leq C_2 \|\xi\|_{W^{1-1/(2r), 2r}(\partial\Omega_0)}^2, \quad (15)$$

for all $\xi \in \Theta$.

iii) Assume Ω_0 is semi-convex and L is of the form (7). Then (14), (15) hold for all $r \in (1, \infty)$.

iv) Assume Ω_0 is semi-convex, L of the form (7) with $L^* = L$ and E is exactly its associated energy (meaning that $\partial_U K(\cdot, U, \nabla U) - \nabla \cdot \partial_q K(\cdot, U, \nabla U) = LU$, for every smooth U). Then (14), (15) hold for all $r \in [1, \infty)$.

Proof. This result is a consequence of Theorem 3.13 which provides an estimate in terms of “interior” norms of ξ , which after replacing $p \in (2, \bar{p}_1)$ by $r = p/(p-2) \in (\bar{r}_1 := \bar{p}_1/(\bar{p}_1-2), \infty)$ gives

$$|E'(\Omega_0)(\xi)| \leq C_1 \|\xi\|_{W^{1, r}(\Omega_0)}, \quad |E''(\Omega_0)(\xi, \xi)| \leq C_2 \|\xi\|_{W^{1, 2r}(\Omega_0)}^2. \quad (16)$$

If Ω_0 is semi-convex then we can have any $p \in (2, \infty)$ which leads to any $r \in (1, \infty)$, and if moreover L is self-adjoint and K is the associated energy, then we can have $p = \infty, r = 1$.

Now, we use the well-known fact, that $E'(\Omega_0)\xi, E''(\Omega_0)(\xi, \xi)$ depend only on the values of ξ on $\partial\Omega_0$. More precisely,

$$\xi, \hat{\xi} \in W^{1, \infty}(B_R), \quad \xi - \hat{\xi} \in W_0^{1, 2r}(\Omega_0) \Rightarrow E'(\Omega_0)(\xi) = E'(\Omega_0)(\hat{\xi}), \quad E''(\Omega_0)(\xi, \xi) = E''(\Omega_0)(\hat{\xi}, \hat{\xi}).$$

Indeed, let ζ_n be a sequence in $C_0^\infty(\Omega_0, \mathbb{R}^N)$ converging to $\xi - \hat{\xi}$ in $W^{1, 2r}(\Omega_0)$. Let $z_n \in C^1([0, \infty) : \mathbb{R}^N)$ be the solution of

$$\forall t \geq 0, \forall x \in \mathbb{R}^N, \quad \partial_t z_n(t, x) = \zeta_n(z_n(t, x)), \quad z_n(0, x) = x.$$

Note that $z_n(t, \cdot) \in \Theta$ (see [17]), and for x in some neighborhood of $\partial\Omega_0$, $z_n(t, x) = x$ for all $t \geq 0$. It follows that $z_n(t, \Omega_0) = \Omega_0$, where $z_n(t, \Omega_0) = \{z_n(t, x), x \in \Omega_0\}$, and therefore $E(z_n(t, \Omega_0)) = E(\Omega_0)$ for all $t \geq 0$. In particular

$$\begin{aligned} E'(\Omega_0)(\xi - \hat{\xi}) &= \lim_{n \rightarrow \infty} \frac{d}{dt} \Big|_{t=0} E(z_n(t, \Omega_0)) = 0, \\ E''(\Omega_0)(\xi - \hat{\xi}, \xi - \hat{\xi}) &= \lim_{n \rightarrow \infty} \frac{d^2}{dt^2} \Big|_{t=0} E(z_n(t, \Omega_0)) = 0. \end{aligned}$$

From this property and (16), we deduce

$$|E'(\Omega_0)(\xi)| \leq C_1 \inf \{ \|\hat{\xi}\|_{W^{1, r}(\Omega_0)} : \hat{\xi} \in W^{1, \infty}(\Omega_0; \mathbb{R}^N), \hat{\xi} = \xi \text{ on } \partial\Omega_0 \},$$

and the similar property for $E''(\Omega_0)(\xi, \xi)$ with the $W^{1, 2r}$ -norm. We then apply Lemma 3.1 to deduce (14), (15). \square

Remark 2.4 The result i) of Theorem 2.3 a priori implies that $E'(\Omega_0)$ and $E''(\Omega_0)$ are continuous on Θ for its $W^{1, \infty}(\Omega_0)$ -norm (as a linear and a bilinear form respectively). Using the previous trace analysis, we deduce that they are also continuous for the $W^{1, \infty}(\partial\Omega_0)$ -norm. The estimates (14), (15) improve this a priori information and show that $E'(\Omega_0), E''(\Omega_0)$ are continuous in $W^{1-1/r, r}(\partial\Omega_0)$ and $W^{1-1/(2r), 2r}(\partial\Omega_0)$ respectively: note that the regularity gets stronger as r gets smaller.

In case iv), the estimate (15) is valid with $r = 1$ (which yields the $H^{1/2}$ -continuity) and cannot be improved since, as already noticed in the introduction (see (3)), $E''(\Omega_0)$ has $H^{1/2}$ -coercivity properties.

Again in case iv), it is interesting to notice that (14) with $r = 1$ can be stated as $E'(\Omega_0) \in (L^1(\partial\Omega_0))' = L^\infty(\partial\Omega_0)$. Again this is sharp in general; if for example E is the Dirichlet energy,

$$E(\Omega) = \frac{1}{2} \int_{\Omega} |\nabla U_{\Omega}|^2 - \int_{\Omega} f U_{\Omega},$$

it is well-known that $E'(\Omega_0) = -\frac{1}{2} |\nabla U_{\Omega_0}|^2|_{\partial\Omega_0}$, which indeed belongs to $L^\infty(\partial\Omega_0)$.

Theorem 2.5 (exterior) *Let Ω_0 be as in (13) and Lipschitz and let $f \in W^{m,\infty}(\mathbb{R}^N)$, $m \in \mathbb{N}^*$, with $\text{supp}(f) \Subset B_R$. For $\theta \in \Theta$, let $U_{\theta} \in H_0^{1,0}(\Omega_{\theta}^e)$ be the solution of (11) in Ω_{θ}^e (see Proposition 3.16) and let $\mathcal{E}(\theta) = E(\Omega_{\theta})$ with E given by (10). Then \mathcal{E} is of class C^m near $\theta = 0 \in \Theta$. Moreover, there exists $\bar{r}_1 = \bar{r}_1(\Omega_0)$ satisfying $\bar{r}_1 \in (1, 2)$ if $N = 2$ and $\bar{r}_1 \in (1, 3)$ if $N \geq 3$ such that, for any $r \in (\bar{r}_1, \infty)$, there exist $C_i = C_i(\Omega_0, U_0, R, f, r)$ with*

$$|E'(\Omega_0)(\xi)| \leq C_1 \|\xi\|_{W^{1-1/r,r}(\partial\Omega_0)} \text{ if } m \geq 1, \quad (17)$$

$$|E''(\Omega_0)(\xi, \xi)| \leq C_2 \|\xi\|_{W^{1-1/(2r),2r}(\partial\Omega_0)}^2 \text{ if } m \geq 2, \quad (18)$$

for any $\xi \in \Theta$.

Proof. Similarly to the proof of Theorem 2.3, this result is a consequence of Theorem 3.19, combined with the trace result of Lemma 3.1. \square

Previous estimates can be written using a different range of Sobolev spaces. In the following statement, we write these estimates in the H^s -norms, which are relevant for our applications (see Section 4).

Corollary 2.6 *Let Ω_0 be as in (13).*

i) *Under the same conditions as in ii) of Theorem 2.3, or as in Theorem 2.5, and with $\bar{r}_1 = \bar{r}_1(\Omega_0)$ as in these theorems, we have*

$$|E'(\Omega_0)(\xi)| \leq C_1 \|\xi\|_{H^s(\partial\Omega_0)}, \quad \forall s \in (s_1, 1], \quad s_1 = \min \left\{ 1, 1 - \frac{1}{\bar{r}_1} + (N-1) \left(\frac{1}{2} - \frac{1}{\bar{r}_1} \right)^+ \right\}, \quad (19)$$

$$|E''(\Omega_0)(\xi, \xi)| \leq C_2 \|\xi\|_{H^s(\partial\Omega_0)}^2, \quad \forall s \in (s_2, 1], \quad s_2 = \min \left\{ 1, \frac{N+1}{2} - \frac{N}{2\bar{r}_1} \right\}, \quad (20)$$

for every $\xi \in \Theta$.

ii) *Under the same conditions as in iii) of Theorem 2.3, (19) (resp. (20)) holds for every $s \in (0, 1]$ (resp. $s \in (1/2, 1]$).*

iii) *Under the same conditions as in iv) of Theorem 2.3, (19) (resp. (20)) holds also for $s = 0$ (resp. $s = 1/2$).*

Proof. Inequalities (19), (20) follow from (14) and (17) respectively and from the embedding result (21) recalled below which may be obtained from [31, Th. 1.107]. Indeed, $\partial\Omega_0$ is an $(N-1)$ -dimensional Lipschitz manifold and since we deal with exponents less than 1, these embeddings carry over to Lipschitz manifolds.

$$\forall t_1, t_2 \in [0, 1], \quad \forall (p_1, p_2) \in [1, \infty],$$

$$\left[t_1 - \frac{N-1}{p_1} > t_2 - \frac{N-1}{p_2} \text{ and } t_1 > t_2 \right] \implies W^{t_1, p_1}(\partial\Omega_0) \subset W^{t_2, p_2}(\partial\Omega_0). \quad (21)$$

We apply this result with $t_1 = s \in [0, 1], p_1 = 2$ and, $[t_2 = 1 - 1/r, p_2 = r]$ to obtain (19) and $[t_2 = 1 - 1/(2r), p_2 = 2r]$ to obtain (20).

For the proof of ii) and iii) we apply the same embeddings and iii), iv) from Theorem 2.3. \square

Remark 2.7 The restrictions on s_1, s_2 in the Lipschitz case i) of Corollary 2.6 are coming from the restriction $p < \bar{p}_1 = \bar{p}_1(\Omega_0)$ as indicated in Theorems 3.9, 3.18. We do not know the exact value of this \bar{p}_1 , but we know that $\bar{p}_1 > 4$ if $N = 2$ and $\bar{p}_1 > 3$ if $N \geq 3$. Thus (recall that $\bar{r}_1 = \bar{p}_1/(\bar{p}_1 - 2)$), we may write $\bar{r}_1 = 2 - \sigma$ if $N = 2$ and $\bar{r}_1 = 3 - \sigma$ if $N \geq 3$, for some small $\sigma > 0$.

In the range of H^s -spaces, the best estimates we obtain in the case i) of Corollary 2.6 may also be written as follows where the constants C_1, C_2 here do not depend on $\xi \in \Theta$, but may depend on other variables, especially ε):

- if $N = 2$, then there exists $\varepsilon > 0$ small such that

$$|E'(\Omega_0)(\xi)| \leq C_1 \|\xi\|_{H^{1/2-\varepsilon}(\partial\Omega_0)}, \quad (22)$$

$$|E''(\Omega_0)(\xi, \xi)| \leq C_2 \|\xi\|_{H^{1-\varepsilon}(\partial\Omega_0)}^2. \quad (23)$$

- If $N = 3$, then there exists $\varepsilon > 0$ small such that

$$|E'(\Omega_0)(\xi)| \leq C_1 \|\xi\|_{H^{1-\varepsilon}(\partial\Omega_0)}.$$

3 Estimates of shape derivatives

3.1 Description of the method

We describe in this paragraph the main idea of the shape derivative estimates on the model example

$$E(\Omega) = \int_{\Omega} K(U_{\Omega}, \nabla U_{\Omega}) := \int_{\Omega} \frac{1}{2} |\nabla U_{\Omega}|^2,$$

where U_{Ω} is solution of (7) with $L = -\Delta$ and Ω, Ω_0 are as in (13). The main point of our approach is that we will make *interior estimates* involving $W^{1,q}(\Omega_0)$ -norms of the directions $\xi \in \Theta$ of differentiation. As already explained in the proof of Theorem 2.3, we will then use the trace Lemma 3.1 to obtain estimates in terms of $W^{1-1/q,q}(\partial\Omega_0)$ -norms of ξ . By doing so, we avoid integrations by parts which would be impossible due to the poor regularity of the boundary of Ω_0 . Moreover, as we will see below, the shape derivative estimates will only depend on the regularity of $U_0 := U_{\Omega_0}$.

One first has to show that E is differentiable.

By changing variable $x = (I + \theta)(y)$, for $\theta \in \Theta$ small, we have

$$\mathcal{E}(\theta) = \int_{\Omega_0} \frac{1}{2} \nabla \hat{U}_{\theta} \cdot \hat{M}_{\theta} \cdot \nabla \hat{U}_{\theta} dx,$$

where

$$\hat{U}_{\theta} = U_{\theta} \circ (I + \theta), \quad \hat{M}_{\theta} = [I + \nabla\theta]^{-1} \mathfrak{t}[I + \nabla\theta]^{-1} \det[I + \nabla\theta], \quad U_{\theta} = U_{\Omega_{\theta}}. \quad (24)$$

Note that $\theta \mapsto \hat{M}_{\theta}$ is C^{∞} from Θ to $(L^{\infty}(\Omega_0))^{N \times N}$ near $\theta = 0$, see [12, 29]. Then the differentiability of $\theta \mapsto \mathcal{E}(\theta)$ fully depends on the differentiability of \hat{U}_{θ} . As it is classical, under reasonable assumptions on L and the data, by using the implicit function theorem, one can prove that $\theta \in \Theta \mapsto \hat{U}_{\theta} \in H_0^1(\Omega_0)$ is of class

C^m near $\theta = 0$, which implies that \mathcal{E} is of class C^m near $\theta = 0$.

Next, let us fix a direction $\xi \in \Theta$; we will denote by $(\cdot)'$ the derivatives with respect to θ in the direction ξ . In general, U_0 is more regular than $H^1(\Omega_0)$ and we consider $p \in (2, \infty)$ such that $U_0 \in W^{1,p}(\Omega_0)$ (see Propositions 3.3 and 3.5 below). We may write:

$$\mathcal{E}'(\theta)(\xi) = \int_{\Omega_0} \nabla \hat{U}_\theta \cdot \hat{M}_\theta \cdot \nabla \hat{U}'_\theta + \frac{1}{2} \nabla \hat{U}_\theta \cdot \hat{M}'_\theta \cdot \nabla \hat{U}_\theta, \quad (25)$$

which at $\theta = 0$ implies (see(24))

$$\begin{aligned} |\mathcal{E}'(0)(\xi)| &\leq C \left(\left| \int_{\Omega_0} \nabla U_0 \cdot \nabla \hat{U}'_0 \right| + \int_{\Omega_0} |\nabla U_0|^2 |\nabla \xi| \right) \\ &\leq C \left(\left| \int_{\Omega_0} \nabla U_0 \cdot \nabla \hat{U}'_0 \right| + \|U_0\|_{W^{1,p}(\Omega_0)}^2 \|\xi\|_{W^{1,p/(p-2)}(\Omega_0)} \right). \end{aligned} \quad (26)$$

The second term in the last estimate being satisfactory, it remains to estimate the first term which involves \hat{U}'_0 . Note first that it would not help much to use a too simple Hölder inequality like

$$\left| \int_{\Omega_0} \nabla U_0 \cdot \nabla \hat{U}'_0 \right| \leq C \|U_0\|_{H^1(\Omega_0)} \|\xi\|_{W^{1,\infty}(\Omega_0)},$$

which uses only the starting information that \hat{U}'_0 is a linear continuous map from Θ to $H^1(\Omega_0)$.

Nor is it appropriate to write the term \hat{U}'_0 in terms of the shape derivative U'_0 , i.e. $\hat{U}'_0 = U'_0 + \nabla U_0 \cdot \xi$. Indeed, in such a case one would have

$$\int_{\Omega_0} \nabla U_0 \cdot \nabla \hat{U}'_0 = \int_{\Omega_0} \nabla U_0 \cdot \nabla U'_0 + \nabla U_0 \cdot D^2 U_0 \cdot \xi + \nabla U_0 \cdot \nabla \xi \cdot \nabla U_0.$$

But we would need here regularity for $D^2 U_0$ and it is not available for the case we are interested in. Even if in the semi-convex situation, we can get some significant information on the first derivative, it becomes quite more difficult for the second derivative (see however [20] for some progress in this direction).

For these reasons, we proceed with the estimate of the term with \hat{U}'_0 by going back the state equation : $LU_\theta = f$, $U_\theta \in H_0^1(\Omega_\theta)$. Recall that we chose $L = -\Delta$ for simplicity here (but the ideas will be the same for general L). The weak form of $LU_\theta = f$ in Ω_θ transported in Ω_0 is

$$\int_{\Omega_0} \varphi \hat{L}_\theta \hat{U}_\theta dx = \int_{\Omega_0} \hat{f}_\theta \varphi dx, \quad \text{for all } \varphi \in H_0^1(\Omega_0), \quad (27)$$

with $\hat{f}_\theta = f \circ (I + \theta) \det[I + \nabla \theta]$ and

$$\hat{L}_\theta : H_0^1(\Omega_0) \mapsto H^{-1}(\Omega_0), \quad \hat{L}_\theta \phi = \nabla \cdot (\hat{M}_\theta \cdot \nabla \phi), \quad (28)$$

where \hat{M}_θ is defined in (24). Note that $\hat{L}_0 = L$ and $(\hat{L}_\theta \hat{U}_\theta)' = \hat{L}_\theta \hat{U}'_\theta + \hat{L}'_\theta \hat{U}_\theta$. By differentiating (27) with respect to θ in the direction ξ , we obtain

$$\int_{\Omega_0} \varphi \hat{L}_\theta \hat{U}'_\theta dx = \int_{\Omega_0} \varphi (\hat{f}'_\theta - \hat{L}'_\theta \hat{U}_\theta), \quad (29)$$

which at $\theta = 0$ gives

$$\int_{\Omega_0} \nabla \hat{U}'_0 \cdot \nabla \varphi = \int_{\Omega_0} \varphi (\hat{f}'_0 - \nabla \cdot (\hat{M}'_0 \cdot \nabla U_0)), \quad (30)$$

The estimate (26) suggests to take $\varphi = U_0 \in H_0^1(\Omega_0)$ in (30), which then yields

$$\begin{aligned} \left| \int_{\Omega_0} \nabla U_0 \cdot \nabla \hat{U}'_0 \right| &\leq C(\Omega_0, f) \int_{\Omega_0} (|U_0| + |\nabla U_0|^2)(|\xi| + |\nabla \xi|) \\ &\leq C(\Omega_0, f, p)(1 + \|U_0\|_{W^{1,p}(\Omega_0)}) \|U_0\|_{W^{1,p}(\Omega_0)} \|\xi\|_{W^{1,p/(p-2)}(\Omega_0)}. \end{aligned} \quad (31)$$

Thus, going back to (26), we deduce

$$|\mathcal{E}'(0)(\xi)| \leq C(\Omega_0, U_0, f, p) \|\xi\|_{W^{1,p/(p-2)}(\Omega_0)}. \quad (32)$$

For more general K and L , the choice of φ is more involved (see the proof of Theorem 3.9), but the procedure is the same.

In the same spirit, that is to say by differentiating (25) and (29) at $\theta = 0$, we can obtain a similar estimate for $|\mathcal{E}''(0)(\xi, \xi)|$. For more details see Sections 3.3, 3.4.

3.2 The trace lemma

Lemma 3.1 *Let Ω be as in (13) and Lipschitz. Let also $q \in [1, \infty]$, $s \in \left(\frac{1}{q}, 1 + \frac{1}{q}\right)$, or $q = s = 1$. Then there exists $C = C(s, q, \Omega)$ such that, for every $\xi \in W^{1,\infty}(\Omega, \mathbb{R}^N) \cap W^{s,q}(\Omega, \mathbb{R}^N)$,*

$$\inf \left\{ \|\hat{\xi}\|_{W^{s,q}(\Omega)}, \hat{\xi} \in W^{1,\infty}(\Omega, \mathbb{R}^N), \hat{\xi} = \xi \text{ on } \partial\Omega \right\} \leq C \|\xi\|_{W^{s-\frac{1}{q},q}(\partial\Omega)}. \quad (33)$$

The main tool for the proof of this lemma is the following classical trace/extension Theorem:

Theorem 3.2 *Let Ω be a set with Lipschitz boundary, $q \in [1, \infty]$ and $s \in \left(\frac{1}{q}, 1 + \frac{1}{q}\right)$ or $q = s = 1$. The trace mapping Tr initially defined on $C^\infty(\bar{\Omega})$ extends as a bounded operator from $W^{s,q}(\Omega)$ to $W^{s-\frac{1}{q},q}(\partial\Omega)$. Moreover there exists a bounded linear operator*

$$\text{Ext} : W^{s-\frac{1}{q},q}(\partial\Omega) \rightarrow W^{s,q}(\Omega)$$

such that $\text{Tr} \circ \text{Ext} = \text{Id}$ on $W^{s,q}(\partial\Omega)$. In all cases, Tr operator is linear, while Ext is linear only in the case $p > 1$.

See [9] for the case $q = s = 1$, and for example [15, Th1 p197] for the other cases.

Proof of Lemma 3.1: Let $\xi \in W^{1,\infty}(\Omega, \mathbb{R}^N) \cap W^{s,q}(\Omega, \mathbb{R}^N)$ (we drop the notation \mathbb{R}^N in the following). We remark that $\xi|_{\partial\Omega} \in W^{s-\frac{1}{q},q}(\partial\Omega)$. Using the extension defined in Theorem 3.2, we can define $\hat{\xi} := \text{Ext}(\text{Tr } \xi) \in W^{s,q}(\Omega)$ satisfying

$$\|\hat{\xi}\|_{W^{s,q}(\Omega)} \leq C \|\xi\|_{W^{s-\frac{1}{q},q}(\partial\Omega)}, \quad (34)$$

where $C = C(s, q, \Omega)$.

Therefore $\hat{\xi} - \xi \in W_0^{s,q}(\Omega)$ and is therefore the limit in $W^{s,q}(\Omega)$ of functions $\alpha_n \in C_0^\infty(\Omega)$. In other words, $\hat{\xi}$ is the limit in $W^{s,q}(\Omega)$ of $\varphi_n = \xi + \alpha_n$. Clearly $\varphi_n \in W^{1,\infty}(\Omega) \cap W^{s,q}(\Omega)$ and $\text{Tr}(\varphi_n) = \text{Tr}(\xi)$, so that

$$\inf \left\{ \|\hat{\xi}\|_{W^{s,q}(\Omega)}, \hat{\xi} \in W^{1,\infty}(\Omega) \cap W^{s,q}(\Omega), \hat{\xi} = \xi \text{ on } \partial\Omega \right\} \leq \|\varphi_n\|_{W^{s,q}(\Omega)},$$

and letting $n \rightarrow \infty$ and using (34), we obtain the first estimate (33). \square

3.3 Shape derivative estimates for an interior PDE

In this section we will prove estimates for the first and second order shape derivatives of the energy (5). As explained above, it will rely on estimates of U_Ω .

3.3.1 Existence, uniqueness and regularity of the solution U_Ω

Proposition 3.3 *Let Ω be as in (13) and Lipschitz. Let L be as in (7) with the matrix (a_{ij}) satisfying (8),(9).*

- i) *Assume $b_i, c \in L^\infty(\Omega)$ and $c \geq 0$. If $f \in H^{-1}(\Omega)$, then there exists a unique solution $U_\Omega \in H_0^1(\Omega)$ of (7).*
- ii) *Assume $a_{ij} \in W^{1,\infty}(\Omega)$, $b_i, c \in L^\infty(\Omega)$ and $c \geq 0$. Then there exists $p_1 = p_1(\Omega)$ satisfying $[p_1 > 4$ if $N = 2$ and $p_1 > 3$ if $N \geq 3]$, such that for every $p \in (p_1', p_1)$ (where $1/p_1 + 1/p_1' = 1$) and every $f \in W^{-1,p}(\Omega)$ the problem (7) admits a unique solution $U_\Omega \in W_0^{1,p}(\Omega)$ and*

$$\|U_\Omega\|_{W^{1,p}(\Omega)} \leq C(\Omega)\|f\|_{W^{-1,p}(\Omega)}. \quad (35)$$

Proof. The point i) is standard, see [8]. The sharp regularity result in ii) has been first proved for $L = -\Delta$ in [13, Thm. 0.5]. The complete proof of ii) is based on [30, Theorem C] and Remark 3.6 for $Lu = -\sum_{i,j} \partial_i(a_{ij}\partial_j u)$, and Remark 3.7 for L of general form. See also Remark 3.8 for a different proof with a slightly stronger regularity on a_{ij} . Note that, according to [30], the same result is valid with quite weaker regularity on a_{ij} , like asking that they be in $\text{VMO}(\mathbb{R}^N)$. \square

Remark 3.4 Notation: If L satisfies the assumptions of ii) above, then so does its adjoint L^* as we can easily see by writing $(b_i u)_{x_i} = (b_i)_{x_i} u + b_i u_{x_i}$, where $b_i \in W^{1,\infty}(\Omega)$. Let p_1, p_1' (resp. $p_1^*, p_1'^*$) be the numbers associated to the equation $LU = f, U \in W_0^{1,p}(\Omega)$ (resp. $L^*U = f, U \in W_0^{1,p}(\Omega)$) as given by ii), Proposition 3.3. Then we set

$$\bar{p}_1 := \min\{p_1, p_1^*\}, \quad \bar{p}_1' := \max\{p_1', p_1'^*\}.$$

Note that, if Ω is Lipschitz, then $\bar{p}_1 > 4$ if $N = 2$ and $\bar{p}_1 > 3$ if $N \geq 3$.

Proposition 3.5 *Let Ω be as in (13) and semi-convex. Let L be as in (7) with the matrix (a_{ij}) satisfying (8),(9).*

- i) *Assume $a_{ij} \in W^{1,\infty}(\Omega)$, $b_i, c \in L^\infty(\Omega)$ and $c \geq 0$. Then for every $p \in (1, \infty)$ and for all $f \in W^{-1,p}(\Omega)$, the problem (7) admits a unique solution $U_\Omega \in W_0^{1,p}(\Omega)$ and (35) holds.*
- ii) *Assume $a_{ij} \in W^{1,\infty}(\Omega)$, $b_i, c \in L^\infty(\Omega)$ and $c \geq 0$. Then for every $f \in L^\infty(\Omega)$ there exists a unique solution $U_\Omega \in H_0^1(\Omega) \cap W^{1,\infty}(\Omega)$ of (7).*

Proof. The result in i) was proven for $L = -\Delta$, see [6, Corollary 1 and Remark]. For general L , we refer to [14] as explained in Remark 2.2 and we use Remark 3.7 below to include the first order terms. For ii), as $f \in L^\infty(\Omega)$, from i) it follows that $U_\Omega \in W^{1,p}(\Omega)$ for all $p \in (1, \infty)$, which implies $U_\Omega \in C^0(\bar{\Omega})$. Furthermore, applying [8, Thm. 8.8], it follows $u \in W_{loc}^{2,2}(\Omega)$. Applying [8, Theorem 9.13] (locally in Ω , with $T = \emptyset$) and then [8, Lemma 9.16], we get $u \in W_{loc}^{2,q}(\Omega)$, for all $q > 2$. From Sobolev embeddings it follows $u \in C^0(\bar{\Omega}) \cap C^2(\Omega)$. Then ii) follows from [8, Thm. 15.9]. \square

Remark 3.6 Theorem C in [30] states that if $a_{ij} \in \text{VMO}(\mathbb{R}^N)$ (and in particular if $a_{ij} \in W^{1,\infty}(\Omega)$), and $L = -\partial_i(a_{ij}(x)\partial_j)$, then $T := \nabla(L)^{-1/2}$ is continuous from $L^p(\Omega)$ to $(L^p(\Omega))^N$, for $p \in (1, p_1)$, with $p_1 = p_1(N)$, $p_1(2) > 4$, $p_1(N) > 3$ for $N \geq 3$. Then it follows that L is continuous invertible from $W_0^{1,p}(\Omega)$ onto $W^{-1,p}(\Omega)$ for $p \in (p'_1, p_1)$. Indeed, first note that this is equivalent to show that $\nabla(L)^{-1}\text{div}$ is continuous from $(L^p(\Omega))^N$ to $L^p(\Omega)$. Next, we note that from [30, Theorem C], $T^* = (L)^{-1/2}\text{div}$ is continuous from $(L^q(\Omega))^N$ to $L^q(\Omega)$, for $q \in (p'_1, \infty)$. Then the desired continuity for $\nabla(L)^{-1}\text{div}$ follows from the fact that $\nabla(L)^{-1}\text{div} = TT^*$.

Remark 3.7 To complete the proof of ii) in Proposition 3.3 and of i) in Proposition 3.5, let us check that if the map $u \rightarrow Au := -\partial_i(a_{ij}(x)\partial_j u)$ defines an isomorphism from $W_0^{1,p}(\Omega)$ into $W^{-1,p}(\Omega)$, then so does L . Indeed, let $B = L - A$, $Bu = b_i\partial_i u + cu$, and consider the equation

$$Lu = f, \quad u \in W_0^{1,p}(\Omega), \quad f \in W^{-1,p}(\Omega), \quad (36)$$

which is equivalent to $Au + Bu = f$. Multiplying this equation by A^{-1} , we get the equivalent equation

$$(I + K)u = g, \quad g = A^{-1}f \in W_0^{1,p}(\Omega), \quad u \in W_0^{1,p}(\Omega), \quad (37)$$

where $K = A^{-1}B : W_0^{1,p}(\Omega) \mapsto W_0^{1,p}(\Omega)$ is compact since B is compact from $W_0^{1,p}$ into $W^{-1,p}$ and A^{-1} is continuous from $W^{-1,p}$ into $W_0^{1,p}$.

Then $I + K$ satisfies the Fredholm alternative (see e.g. [4, Thm. 6.6]). Furthermore, $\text{Ker}(I + K) = \text{Ker}(L) = \{0\}$. Indeed, assume $u \in W_0^{1,p}(\Omega)$, $Lu = 0$. If $p \geq 2$, we can directly use the uniqueness result in [8, Thm. 8.1 and Cor. 8.2] to deduce $u = 0$. If $p \in (1, 2)$, we obtain the same conclusion by using that $L^* : W_0^{1,p'} \mapsto W^{-1,p'}$ is onto which follows from [8, Thm.8.3 and 8.6] (see also the remark following Corollary 8.7 in [8]).

Then, Fredholm alternative [4, Thm. 6.6] implies that $I + K$ defines an isomorphism from $W_0^{1,p}(\Omega)$ onto itself. \square

Remark 3.8 Different approaches may be quoted for the last point of Proposition 3.3. They only require a slightly stronger regularity hypothesis (like $a_{ij} \in C^{1+\gamma}(\Omega)$, $\gamma \in (0, 1)$), but they are also quite interesting. Let us just recall the idea: we need to focus only in the case $L = -\partial_i(a_{ij}(x)\partial_j)$, because the case of general L follows from Remark 3.7. The case $N = 2$ follows from [22]. For the case $N \geq 3$ we consider the manifold (Ω, g) , where g is the metric given by $g = \det(a_{ij})^{1/(N-2)}A^{-1}$. Then (from [25, p. 186] and the formula of Laplace-Beltrami operator, see [22, (1.1)] for instance), the equation $Lu = f$, $u \in W_0^{1,p}(\Omega)$ with $f \in W^{-1,p}(\Omega)$, becomes $-\Delta_g u = h$, with $h = \det(a_{ij})^{1/(2-N)}f \in W^{-1,p}(\Omega)$. Note that $g \in C^{1,1}(\Omega)$ and $W^{-1,p}(\Omega)$ norms of f and h are equivalent. Then the claim ii) follows from [24, Corollary 13.2].

3.3.2 Estimates of the shape derivatives of the solution

The results of this section give estimates for \hat{U}'_0 and \hat{U}''_0 . Note that while for \hat{U}'_0 we prove an $H^1(\Omega_0)$ -estimate, we also prove more involved estimates for \hat{U}'_0 and \hat{U}''_0 . These estimates are motivated by the functional K involved in the energy (5), see also Theorem 3.13.

Theorem 3.9 *Let Ω_0 be as in (13) and let L be as in (7) with the matrix (a_{ij}) satisfying (8),(9). Let $\theta \in \Theta$, $\Omega_\theta = (I + \theta)(\Omega_0)$, $U_\theta \in H_0^1(\Omega_\theta)$ solution of (7) (see i) in Proposition 3.3), $\hat{U}_\theta = U_\theta \circ (I + \theta)$.*

- i) *If $a_{ij}, b_i, c, f \in W^{m,\infty}(B_R)$, $m \in \mathbb{N}^*$, then $\hat{U}_\theta \in C^m(\mathcal{V}, H_0^1(\Omega_0))$ where \mathcal{V} is a neighborhood of $\theta = 0$ in Θ .*

In the following, we assume the previous hypotheses are satisfied for $m = 2$.

ii) If moreover Ω_0 is Lipschitz and $\alpha_{ij} \in W^{1,\infty}(\Omega)$, $\beta_i, \gamma, \alpha_{00} \in L^\infty(\Omega)$ then for all $p \in (2, \bar{p}_1)$ we have

$$\|\hat{U}'_0\|_{H^1(\Omega_0)} \leq C_1 \|\xi\|_{W^{1,2p/(p-2)}(\Omega_0)}, \quad (38)$$

$$\left| \int_{\Omega_0} \partial_U K(\cdot, U_0, \nabla U_0) \hat{U}'_0 + \partial_q K(\cdot, U_0, \nabla U_0) \cdot \nabla \hat{U}'_0 \right| \leq C_1 \|\xi\|_{W^{1,p/(p-2)}(\Omega_0)}, \quad (39)$$

$$\left| \int_{\Omega_0} \partial_U K(\cdot, U_0, \nabla U_0) \hat{U}''_0 + \partial_q K(\cdot, U_0, \nabla U_0) \cdot \nabla \hat{U}''_0 \right| \leq C_2 \|\xi\|_{W^{1,2p/(p-2)}(\Omega_0)}^2, \quad (40)$$

where \bar{p}_1 is defined in Remark 3.4 ($\bar{p}_1 > 4$ if $N = 2$, $\bar{p}_1 > 3$ if $N \geq 3$), $\hat{U}'_0 = \hat{U}'_\theta(0)(\xi)$, $\hat{U}''_0 = \hat{U}''_\theta(0)(\xi, \xi)$, $C_i = C_i(\Omega_0, L, f, K, p)$.

Proof.

Step 1. The proof of i) is classical. It is based on the implicit function theorem.

After changing the variable $x = (I + \theta)(y)$, the weak form of (7) in Ω_θ is transformed in Ω_0 as follows

$$\langle \hat{L}_\theta \hat{U}_\theta, \varphi \rangle_{H^{-1}(\Omega_0) \times H_0^1(\Omega_0)} = \int_{\Omega_0} \varphi \hat{f}_\theta, \quad \forall \varphi \in H_0^1(\Omega_0), \quad (41)$$

where $\hat{L}_\theta : H_0^1(\Omega_0) \mapsto H^{-1}(\Omega_0)$ is defined by

$$\begin{aligned} \langle \hat{L}_\theta \hat{U}, \varphi \rangle &= \langle \partial_l(\hat{a}_{kl} \partial_k \hat{U}) + \hat{b}_k \partial_k \hat{U} + \hat{c} \hat{U}, \varphi \rangle \\ &= \int_{\Omega_0} \hat{a}_{kl} \partial_k \hat{U} \partial_l \varphi + (\hat{b}_k \partial_k \hat{U} + \hat{c} \hat{U}) \varphi, \quad \text{for } (\hat{U}, \varphi) \in H_0^1(\Omega_0)^2, \end{aligned} \quad (42)$$

and

$$\begin{aligned} \hat{a}_{kl} &= \sum_{i,j} a_{ij} \circ (I + \theta) T_{ki} T_{lj} \det[I + \nabla \theta], & \hat{b}_k &= \sum_i b_i \circ (I + \theta) T_{ki} \det[I + \nabla \theta], \\ \hat{c} &= c \circ (I + \theta) \det[I + \nabla \theta], & \hat{f} &= f \circ (I + \theta) \det[I + \nabla \theta], \\ T &= (T_{kl}) = [Id + \nabla \theta]^{-1}. \end{aligned}$$

Consider F defined by

$$\begin{aligned} F : \Theta \times H_0^1(\Omega_0) &\longrightarrow H^{-1}(\Omega) \\ (\theta, \hat{U}) &\longmapsto \left[\varphi \mapsto \langle \hat{L}_\theta \hat{U} - \hat{f}, \varphi \rangle \right]. \end{aligned}$$

It is easy to check that F is well defined and of class C^m in a neighborhood of $(0, U_0) \in \Theta \times H_0^1(\Omega_0)$, and

$$\partial_U F(0, U_0)(U) : \varphi \mapsto \langle LU, \varphi \rangle = \int_{\Omega_0} a_{ij} \partial_i U \partial_j \varphi + (b_i \partial_i U + cU) \varphi, \quad \forall U, \varphi \in H_0^1(\Omega_0).$$

Note that from Proposition 3.3, $\partial_U F(0, U_0)$ defines an isomorphism from $H_0^1(\Omega_0)$ to $H^{-1}(\Omega)$. Then, from implicit function Theorem there exists a C^m map, $\hat{U} : \theta \mapsto \hat{U}(\theta)$, such that $F(\theta, \hat{U}(\theta)) = 0$ for $\|\theta\|_\Theta$ small. From the uniqueness of solution to (7) we obtain $\hat{U}_\theta = \hat{U}(\theta)$, which proves the regularity of $\theta \mapsto \hat{U}_\theta$.

Step 2. We prove the estimates (38)-(40) by differentiating (41) with respect to θ . The differentiation is allowed because \hat{U}_θ is differentiable and a_{ij}, b_i, c, f are regular enough.

Differentiating (41) once with respect to θ gives

$$\langle \hat{L}_\theta \hat{U}'_\theta, \varphi \rangle = \langle \hat{f}'_\theta - \hat{L}'_\theta \hat{U}_\theta, \varphi \rangle. \quad (43)$$

We take $\theta = 0$ in (43). Note that $\hat{L}_0 = L$. By isolating all the terms with \hat{U}'_0 we obtain

$$\begin{aligned} \langle L\hat{U}'_0, \varphi \rangle &= \langle L^*\varphi, \hat{U}'_0 \rangle \\ &= \int_{\Omega_0} \hat{f}'_0 \varphi - \left(\hat{a}'_{kl} \partial_k U_0 \partial_l \varphi + (\hat{b}'_k \partial_k U_0 + \hat{c}'_0 U_0) \varphi \right). \end{aligned} \quad (44)$$

We now choose suitable test functions φ to prove (38) and (39). Let $\varphi \in H_0^1(\Omega_0)$ be the solution of

$$L^*\varphi = \hat{U}'_0 - \Delta \hat{U}'_0 =: g^*. \quad (45)$$

Note that $g^* \in H^{-1}(\Omega_0)$ and then the solution φ is uniquely defined in $H_0^1(\Omega_0)$ and satisfies (see [18])

$$\|\varphi\|_{H^1(\Omega_0)} \leq C \|g^*\|_{H^{-1}(\Omega_0)} \leq C \|\hat{U}'_0\|_{H^1(\Omega_0)}, \quad C = C(\Omega_0, L). \quad (46)$$

Then from (44) it follows

$$\begin{aligned} \|\hat{U}'_0\|_{H^1(\Omega_0)}^2 &\leq C_1 \int_{\Omega_0} (1 + |U_0| + |\nabla U_0|)(|\xi| + |\nabla \xi|)(|\varphi| + |\nabla \varphi|) \\ &\leq C_1 (1 + \|U_0\|_{W^{1,p}(\Omega_0)}) \|\xi\|_{W^{1,2p/(p-2)}(\Omega_0)} \|\varphi\|_{H^1(\Omega_0)}, \end{aligned} \quad (47)$$

where we applied Hölder inequality with three terms, and $C_1 = C_1(\Omega_0, L, f)$, $p \in (2, p_1)$, $p_1 = p_1(\Omega_0)$ as given by Proposition 3.3 for L . Together with (46) this proves (38).

Now we consider $\varphi \in H_0^1(\Omega_0)$ the solution of

$$\begin{aligned} L^*\varphi &= \partial_U K(\cdot, U_0, \nabla U_0) - \nabla \cdot \partial_q K(\cdot, U_0, \nabla U_0) \\ &= (\alpha_{00} U_0 + \gamma) - \nabla \cdot (\alpha \nabla U_0 + \beta) =: g^*. \end{aligned} \quad (48)$$

Let $q \in (2, \bar{p}_1)$. Recall that $U_0 \in W^{1,q}(\Omega_0)$ by Proposition 3.3. From (6), we deduce

$$\partial_U K(\cdot, U_0, \nabla U_0), \partial_q K(\cdot, U_0, \nabla U_0) \in L^q(\Omega_0).$$

It follows that $g^* \in W^{-1,q}(\Omega_0)$. By Proposition 3.3 applied to L^* , the solution φ of (48) is uniquely defined in $W_0^{1,q}(\Omega_0)$ and satisfies

$$\|\varphi\|_{W^{1,q}(\Omega_0)} \leq C \|g^*\|_{W^{-1,q}(\Omega_0)} = C, \quad C = C(\Omega_0, L, K, U_0, q). \quad (49)$$

Then from (44) it follows

$$\begin{aligned} \left| \int_{\Omega_0} \partial_U K(\cdot, U_0, \nabla U_0) \hat{U}'_0 + \partial_q K(U_0, \nabla U_0) \cdot \nabla \hat{U}'_0 \right| \\ &= \left| \langle L^*\varphi, \hat{U}'_0 \rangle \right| = \left| \langle L\hat{U}'_0, \varphi \rangle \right| \\ &\leq C_1 \int_{\Omega_0} (1 + |U_0| + |\nabla U_0|)(|\xi| + |\nabla \xi|)(|\varphi| + |\nabla \varphi|) \\ &\leq C_1 (1 + \|U_0\|_{W^{1,p}(\Omega_0)}) \|\varphi\|_{W^{1,q}(\Omega_0)} \|\xi\|_{W^{1,1/(1-1/p-1/q)}(\Omega_0)}, \end{aligned} \quad (50)$$

with $C_1 = C_1(\Omega_0, L, K, f, q)$. Here we chose $q := p$ where p is given in ii) of Theorem 3.9 and the estimate (39) follows. (Note that the use in considering $q \neq p$ in the previous computations will appear later in the case of semi-convex domains).

Step 3. Differentiating (43) at $\theta = 0$ and isolating the terms with \hat{U}''_0 gives

$$\begin{aligned} \langle L\hat{U}''_0, \varphi \rangle &= \int_{\Omega_0} \hat{f}''_0 \varphi - (\hat{a}''_{kl} \partial_k \hat{U}_0 \partial_l \varphi + (\hat{b}''_k \partial_k \hat{U}_0 + \hat{c}''_0 \hat{U}_0) \varphi) \\ &\quad - \int_{\Omega_0} 2 \left(\hat{a}'_{kl} \partial_k \hat{U}'_0 \partial_l \varphi + (\hat{b}'_k \partial_k \hat{U}'_0 + \hat{c}'_0 \hat{U}'_0) \varphi \right). \end{aligned} \quad (51)$$

It implies

$$\left| \langle L^* \varphi, \hat{U}_0'' \rangle \right| \leq C_2 \int_{\Omega_0} \left((1 + |U_0| + |\nabla U_0|)(|\xi|^2 + |\nabla \xi|^2) + (|\hat{U}_0'| + |\nabla \hat{U}_0'|)(|\xi| + |\nabla \xi|) \right) (|\varphi| + |\nabla \varphi|), \quad (52)$$

where $C_2 = C_2(\Omega_0, L, f)$.

Then (52) with φ solution of (48), together with (38) yield

$$\begin{aligned} \left| \langle L^* \varphi, \hat{U}_0'' \rangle \right| &= \left| \int_{\Omega_0} \partial_U K(\cdot, U_0, \nabla U_0) \hat{U}_0'' + \partial_q K(\cdot, U_0, \nabla U_0) \cdot \nabla \hat{U}_0'' \right| \\ &\leq C_2 \left((1 + \|U_0\|_{W^{1,p}(\Omega_0)}) \|\xi\|^2 + \|\nabla \xi\|^2_{L^{1/(1-1/p-1/q)}(\Omega_0)} + \|\nabla \hat{U}_0'\|_{L^2(\Omega_0)} \|\xi\|_{W^{1,1/(1/2-1/q)}(\Omega_0)} \right) \\ &\quad \|\varphi\|_{W^{1,q}(\Omega_0)} \\ &\leq C_2 (1 + \|U_0\|_{W^{1,p}(\Omega_0)}) \|U_0\|_{W^{1,q}(\Omega_0)} \\ &\quad \left(\|\xi\|_{W^{1,2/(1-1/p-1/q)}(\Omega_0)}^2 + \|\xi\|_{W^{1,2p/(p-2)}(\Omega_0)} \|\xi\|_{W^{1,2q/(q-2)}(\Omega_0)} \right), \end{aligned} \quad (53)$$

with $C_2 = C_2(\Omega_0, L, f, K, p, q)$. Again we choose $q := p$ and this proves (40). \square

Proposition 3.10 *Besides the assumptions of ii) in Theorem 3.9, we assume Ω_0 is semi-convex. Then the following holds.*

i) For all $p \in (2, \infty)$ we have

$$\|\hat{U}_0'\|_{H^1(\Omega_0)} \leq C_1 \|\xi\|_{H^1(\Omega_0)}, \quad (54)$$

$$\left| \int_{\Omega_0} \partial_U K(\cdot, U_0, \nabla U_0) \hat{U}_0' + \partial_q K(\cdot, U_0, \nabla U_0) \cdot \nabla \hat{U}_0' \right| \leq C_1 \|\xi\|_{W^{1,p/(p-2)}(\Omega_0)}^2, \quad (55)$$

$$\left| \int_{\Omega_0} \partial_U K(\cdot, U_0, \nabla U_0) \hat{U}_0'' + \partial_q K(\cdot, U_0, \nabla U_0) \cdot \nabla \hat{U}_0'' \right| \leq C_2 \|\xi\|_{W^{1,2p/(p-2)}(\Omega_0)}^2, \quad (56)$$

where $C_i = C_i(\Omega_0, L, f, K, p)$.

ii) If furthermore L is self-adjoint and E is its energy associated, i.e. $\partial_U K(\cdot, U, \nabla U) - \nabla \cdot \partial_q K(\cdot, U, \nabla U) = LU$, then we can take $p = \infty$ in (55), (56).

Proof. We proceed as in Theorem 3.9 using the extra property that Ω_0 is semi-convex. We now have $U_0 \in W^{1,\infty}(\Omega_0)$ (see Proposition 3.5). Moreover, the solution φ of (48) satisfies $\varphi \in W^{1,q}(\Omega)$ for all $q \in (2, \infty)$. Then we can take $p = \infty$ in (47) which implies (54). Furthermore, we can apply (50) and (53) with $p = \infty$ and $q \in (2, \infty)$ arbitrary, which prove (55) and (56).

In the case Ω semi-convex, L self-adjoint and K the energy associated to L , then $L = L^*$ and $L^* \varphi = LU_0$. Therefore, $\varphi = U_0$ in (48). Hence $U_0 \in W^{1,\infty}(\Omega_0)$ and $\varphi \in W^{1,\infty}(\Omega_0)$. Then we proceed as above with $p = q = \infty$. \square

Remark 3.11 The results of Proposition 3.10 hold for any Ω_0 such that $U_0 \in W^{1,\infty}(\Omega_0)$ and such that $\varphi \in W_0^{1,q}(\Omega_0) \mapsto L^* \varphi \in W^{-1,q}(\Omega_0)$ is bounded and invertible for all $q \in (2, \infty)$.

Remark 3.12 In Theorem 3.9, we could try to estimate $\|\hat{U}_0''\|_{H^1(\Omega_0)}$. Indeed, in step 3 of Theorem 3.9, we take φ to be the solution of $L^* \varphi = \hat{U}_0'' - \Delta \hat{U}_0'' =: f^*$, so that $\|\hat{U}_0''\|_{H^1(\Omega_0)}^2 = \langle L^* \varphi, \hat{U}_0'' \rangle$. Then we can proceed

as in (53). However, as we have only $\hat{U}'_0, \hat{U}''_0 \in H^1_0(\Omega_0)$, then $f^* \in H^{-1}(\Omega_0)$ and so $\varphi \in H^1_0(\Omega_0)$ (sharp in general). Therefore, in (53) we have $q = 2$ and it leads to

$$\begin{aligned} \|\hat{U}''_0\|_{H^1(\Omega_0)}^2 &\leq C_2(1 + \|U_0\|_{W^{1,p}(\Omega_0)})\|U_0\|_{W^{1,2}(\Omega_0)} \\ &\quad \left(\|\xi\|_{W^{1,4p/(p-2)}(\Omega_0)}^2 + \|\xi\|_{W^{1,2p/(p-2)}(\Omega_0)}\|\xi\|_{W^{1,\infty}(\Omega_0)} \right), \end{aligned} \quad (57)$$

which is not appropriate as it contains the strong norm $\|\xi\|_{W^{1,\infty}(\Omega_0)}$.

3.3.3 Estimates of the shape derivatives of the energy

Theorem 3.13 *Let Ω_0 be as in (13) and let L be as in (7) with the matrix (a_{ij}) satisfying (8),(9). The following properties hold.*

i) If $a_{ij}, b_i, c, f, \alpha_{ij}, \beta_i, \gamma, \delta \in W^{m,\infty}(B_R)$, $m \geq 1$, then $\mathcal{E} : \theta \mapsto \mathcal{E}(\theta)$ is of class C^m in a neighborhood of $\theta = 0$ in Θ .

In the following, we assume the previous hypotheses are satisfied for $m = 2$.

ii) If Ω_0 is Lipschitz, then for all $p \in (2, \bar{p}_1)$ we have

$$|\mathcal{E}'(0)(\xi)| \leq C_1 \|\xi\|_{W^{1,p/(p-2)}(\Omega_0)}, \quad (58)$$

$$|\mathcal{E}''(0)(\xi, \xi)| \leq C_2 \|\xi\|_{W^{1,2p/(p-2)}(\Omega_0)}^2, \quad (59)$$

where $C_i = C_i(\Omega_0, L, f, K, p)$ and \bar{p}_1 (introduced in Remark 3.4) satisfies $\bar{p}_1 > 4$ if $N = 2$ and $\bar{p}_1 > 3$ if $N \geq 3$.

iii) If Ω_0 is semi-convex, then (58) and (59) hold for all $p \in (2, \infty)$.

iv) If Ω_0 is semi-convex, L is self-adjoint and K is its associated energy, then (58) and (59) hold for $p = \infty$.

Remark 3.14 Theorem 3.13 can be stated in a more general form, based on the regularity of the state solution related to the operator L and its adjoint L^* . Namely, if

a) $a_{ij}, b_i, c, f, \alpha_{ij}, \beta_i, \gamma, \delta \in W^{2,\infty}(\Omega_0)$,

b) the operators L and L^* define isomorphisms from $W_0^{1,p}(\Omega) \mapsto W^{-1,p}(\Omega)$, for some $p \in [1, \infty]$

then (58) and (59) hold for this p .

Proof. [of Theorem 3.13]

Step 1. Note that by changing the variable $y = (I + \theta)(x)$ we have

$$\begin{aligned} \mathcal{E}(\theta) &= \int_{\Omega_0} K(I + \theta, \hat{U}_\theta, {}^t[T] \cdot \nabla \hat{U}_\theta) \det[I + \nabla \theta] dx \\ &= \int_{\Omega_0} \left(\frac{1}{2} \hat{\alpha}_{kl} \partial_k \hat{U}_\theta \partial_l \hat{U}_\theta + \frac{1}{2} \hat{\alpha}_{00} \hat{U}_\theta^2 + \hat{\beta}_k \partial_k \hat{U}_\theta + \hat{\gamma} \hat{U}_\theta + \hat{\delta} \right), \end{aligned} \quad (60)$$

where

$$\begin{aligned} \hat{\alpha}_{kl} &= \alpha_{ij} \circ (I + \theta) T_{ki} T_{lj} \det[I + \nabla \theta], & \hat{\beta}_i &= \beta_i \circ (I + \theta) T_{ki} \det[I + \nabla \theta], \\ \hat{\gamma} &= \gamma \circ (I + \theta) \det[I + \nabla \theta], & \hat{\delta} &= \delta \circ (I + \theta) \det[I + \nabla \theta]. \end{aligned}$$

The differentiability of \mathcal{E} follows from the regularity of K and i), Theorem 3.9.

Step 2. Assume Ω_0 is Lipschitz. Differentiating (60) gives

$$\begin{aligned} \mathcal{E}'(\theta)(\xi) &= \int_{\Omega_0} \hat{\alpha}_{kl} \partial_k \hat{U}'_\theta \partial_l \hat{U}_\theta + \hat{\alpha}_{00} \hat{U}_\theta \hat{U}'_\theta + \hat{\beta}_i \partial_k \hat{U}'_\theta + \hat{\gamma} \hat{U}'_\theta \\ &\quad + \int_{\Omega_0} \left(\frac{1}{2} \hat{\alpha}'_{kl} \partial_k \hat{U}_\theta \partial_l \hat{U}_\theta + \frac{1}{2} \hat{\alpha}'_{00} \hat{U}_\theta^2 + \hat{\beta}'_k \partial_k \hat{U}_\theta + \hat{\gamma}' \hat{U}_\theta + \hat{\delta}' \right). \end{aligned} \quad (61)$$

Taking $\theta = 0$ in (61) and using (39) gives

$$\begin{aligned} |\mathcal{E}'(0)(\xi)| &\leq \left| \int_{\Omega_0} \partial_U K(x, U_0, \nabla U_0) \hat{U}'_0 + \partial_q K(x, U_0, \nabla U_0) \cdot \nabla \hat{U}'_0 \right| \\ &\quad + C_1 \int_{\Omega_0} (1 + |U_0|^2 + |\nabla U_0|^2) (|\xi| + |\nabla \xi|) \\ &\leq C_1 (\|\xi\|_{W^{1,q/(q-2)}(\Omega_0)} + (1 + \|U_0\|_{W^{1,p}(\Omega_0)}^2) \|\xi\|_{W^{1,p/(p-2)}(\Omega_0)}), \end{aligned} \quad (62)$$

with $C_1 = C_1(\Omega_0, L, f, K, p)$ and $p, q \in (2, \bar{p}_1)$.

Differentiating (61) at $\theta = 0$, isolating the terms with \hat{U}''_0 , using the $W^{2,\infty}$ -regularity of the coefficients of L and K , the $W^{1,p}(\Omega_0)$ regularity of U_0 and the estimates (38), (40) yields

$$\begin{aligned} |\mathcal{E}''(0)(\xi, \xi)| &\leq \left| \int_{\Omega_0} \partial_U K(x, U_0, \nabla U_0) \hat{U}''_0 + \partial_q K(x, U_0, \nabla U_0) \cdot \nabla \hat{U}''_0 \right| \\ &\quad + C_2 \int_{\Omega_0} \left[|\hat{U}'_0|^2 + |\nabla \hat{U}'_0|^2 + (1 + |U_0|^2 + |\nabla U_0|^2) (|\xi|^2 + |\nabla \xi|^2) \right] \\ &\leq C_2 (\|\xi\|_{W^{1,2q/(q-2)}(\Omega_0)}^2 + (1 + \|U_0\|_{W^{1,p}(\Omega_0)}^2) \|\xi\|_{W^{1,2p/(p-2)}(\Omega_0)}^2), \end{aligned} \quad (63)$$

with $C_2 = C_2(\Omega_0, L, f, K, p)$ and $p, q \in (2, \bar{p}_1)$.

Taking $q = p$ in (62), (63) proves ii).

Step 3. If Ω_0 is semi-convex then from $U_0 \in W^{1,\infty}(\Omega_0)$ and i), Proposition 3.10, we can take $p = \infty$ and $q \in (2, \infty)$ in (62) and (63), which proves iii).

Step 4. Finally, if Ω_0 is semi-convex, $L = L^*$ and K is the energy associated to L then from $U_0 \in W^{1,\infty}(\Omega_0)$ and ii), Proposition 3.10, we can take $q = p = \infty$ in (62) and (63), which proves iv). \square

Remark 3.15 The technique we have used to obtain estimates for $E'(\Omega)$ and $E''(\Omega)$, can be applied to shape functionals E defined as in (5) but involving more general K . Once the differentiability of E is proven, our computations can be used similarly to get estimates whose exponents will depend on the growth of K at infinity. Note that the proof of the differentiability of E may rely in these cases on the differentiability of the map $\theta \in \Theta \mapsto \hat{U}(\theta) \in W^{1,p}(\Omega_0)$ with certain $p \geq 2$, which is not known as far as we know.

3.4 Shape derivative estimates for an exterior PDE

In this section we will apply the technique described in Section 3.1 to estimate the first and second order shape derivatives of the energy (10), related to the problem (11) in the exterior of a domain Ω .

Even in the case when Ω is convex, its exterior $\Omega^e = \mathbb{R}^N \setminus \bar{\Omega}$ is only a Lipschitz domain. Therefore, subject to the existence, uniqueness and regularity of the solution of the problem (11), which represents some particularities as it is in the exterior of Ω , the method we developed in Section 3.1 and used for the interior problem in Section 3.3, also applies for the problem in the exterior.

As in the previous section, we consider $\Theta = W^{1,\infty}(B_R, \mathbb{R}^N)$ and all the domains Ω under consideration will be assumed to satisfy at least (13).

3.4.1 Existence, uniqueness and regularity

To analyze problem (11), it is appropriate to consider the space $H^{1,0}(\Omega)$ as introduced in (12). This space is a Banach space, see [26], and even an Hilbert space if equipped with the inner product

$$(U, V)_{H^{1,0}(\Omega^e)} := \int_{\Omega^e} \left(\frac{UV}{(1 + |x|^2)[\ln(2 + |x|^2)]^{\delta_{2,N}}} + \nabla U \cdot \nabla V \right) dx, \quad (64)$$

where $\delta_{2,N}$ is defined in (12). Let $H_0^{1,0}(\Omega^e)$ denote the closure of $\mathcal{D}(\Omega^e)$ in $H^{1,0}(\Omega^e)$. It can be shown that $H_0^{1,0}(\Omega^e)$ equipped with the inner product

$$(U, V)_{H_0^{1,0}(\Omega^e)} := \int_{\Omega^e} \nabla U \cdot \nabla V dx, \quad (65)$$

is an Hilbert space, see [26]. Furthermore, the norms in $H_0^{1,0}(\Omega^e)$ generated by the inner products (64) and (65) are equivalent. Let $H^{-1,0}(\Omega^e)$ denote the dual space of $H_0^{1,0}(\Omega^e)$

For the solution of (11) we have the following regularity result.

Proposition 3.16 *Let Ω be as in (13). Assume it is also Lipschitz. Let $f \in L^\infty(\mathbb{R}^N)$, $\text{supp}(f) \subset B_R$.*

i) Then there exists a unique weak solution $U \in H_0^{1,0}(\Omega^e)$ to (11) satisfying

$$\|U\|_{H^{1,0}(\Omega^e)} \leq C(\Omega) \|f\|_{H^{-1,0}(\Omega^e)} \leq C(\Omega, R) \|f\|_{L^\infty}. \quad (66)$$

ii) There exists $p_1 = p_1(\Omega_0)$ satisfying [$p_1 > 4$ if $N = 2$ and $p_1 > 3$ if $N \geq 3$], such that $U \in W_0^{1,p}(\Omega^e)$ for every $p \in (p'_1, p_1)$ where $1/p_1 + 1/p'_1 = 1$, and

$$\|U\|_{W^{1,p}(\Omega_R^e)} \leq C(\Omega, R, p) \|f\|_{L^\infty}, \quad \Omega_R^e = \Omega^e \cap B_R. \quad (67)$$

Proof.

Step 1. Note that the weak solution U of (11) satisfies

$$\int_{\Omega^e} \nabla U \cdot \nabla \varphi = \int_{\Omega^e} f \varphi, \quad \forall \varphi \in H_0^{1,0}(\Omega^e). \quad (68)$$

Note also that as $f \in L^\infty(\mathbb{R}^N)$ has compact support, then $f \in L^2(\Omega^e) \subset H^{-1,0}(\Omega^e)$ with $\|f\|_{H^{-1,0}(\Omega^e)} \leq C(R) \|f\|_{L^\infty}$. Therefore existence and uniqueness of a solution $U \in H_0^{1,0}(\Omega^e)$ to (68) together with estimate (66) follows from Lax-Milgram Lemma and from (65).

Step 2. We will use [13, Theorem 0.5]. Let $\eta \in \mathcal{D}(\mathbb{R}^N)$, $0 \leq \eta \leq 1$, $\eta = 1$ in B_R and $\eta = 0$ in $\mathbb{R}^N \setminus B_{2R}$. Then ηU satisfies

$$\begin{aligned} -\Delta(\eta U) &= f\eta - 2\nabla U \cdot \nabla \eta - U\Delta\eta =: g \quad \text{in } \Omega_{2R}^e := \Omega^e \cap B_{2R}, \\ \eta U &= 0 \quad \text{on } \partial\Omega_{2R}^e. \end{aligned}$$

From i) above, $g \in L^2(\Omega_{2R}^e)$. Let $B_{R,2R} := \{x \in \mathbb{R}^N; R < |x| < 2R\}$. From local regularity properties of $-\Delta$, as $-\Delta U = f = 0$ on a neighborhood of the closure of $B_{R,2R}$, we get $U \in W^{1,p}(B_{R,2R})$ for all $p \in (1, \infty)$ with

$$\|U\|_{W^{1,p}(B_{R,2R})} \leq C(\Omega, R, p) \|U\|_{H^{1,0}(\Omega^e)} \leq C(\Omega, R, p) \|f\|_{L^\infty}.$$

Since $\eta = 1$, $\nabla \eta = 0$, $\Delta \eta = 0$ on B_R , it follows that $g \in L^p(\Omega_{2R}^e)$ with $\|g\|_{L^p(\Omega_{2R}^e)} \leq C(\Omega, R, p) \|f\|_{L^\infty}$.

We recall [13, Theorem 0.5] which states that there exists p_1 depending on Ω_{2R}^e , satisfying $p_1 > 4$ if $N = 2$ and $p_1 > 3$ for every $N \geq 3$, such that the map $V \in W_0^{1,p}(\Omega_{2R}^e) \mapsto -\Delta V \in W^{-1,p}(\Omega_{2R}^e)$ defines an isomorphism for every $p \in (p_1', p_1)$ (see also Proposition 3.3). It implies that

$$\|\eta U\|_{W^{1,p}(\Omega_{2R}^e)} \leq C(\Omega, R, p) \|g\|_{W^{-1,p}(\Omega_{2R}^e)} \leq C(\Omega, R, p) \|g\|_{L^p(\Omega_{2R}^e)} \leq C(\Omega, R, p) \|f\|_{L^\infty},$$

which implies (67). \square

Remark 3.17 In connection with Remark 3.4 related to the interior problem, note that with our choice of $L = L^* = -\Delta$, we have here $p_1 = p_1^* = \bar{p}_1$. Recall that $\bar{p}_1 > 4$ if $N = 2$ and $\bar{p}_1 > 3$ if $N \geq 3$.

3.4.2 Estimates of the shape derivatives of the solution

We have the following theorem, which is similar to Theorem 3.9:

Theorem 3.18 *Let Ω_0 be as in (13). Assume it is also Lipschitz. Let $\theta \in \Theta$, $\Omega_0^e = \mathbb{R}^N \setminus \bar{\Omega}_0$, $\Omega_\theta^e = (I + \theta)(\Omega_0^e)$, $f \in L^\infty(\mathbb{R}^N)$ with $\text{supp}(f) \subset B_R$, $U_\theta \in H_0^{1,0}(\Omega_\theta^e)$ the solution of (11) (see i) in Proposition 3.16), $\hat{U}_\theta = U_\theta \circ (I + \theta)$.*

i) *If $f \in W^{m,\infty}(\Omega_{2R}^e)$, $m \in \mathbb{N}^*$, then $\theta \mapsto \hat{U}_\theta \in H_0^1(\Omega_0)$ is of class C^m in a neighborhood of $\theta = 0 \in \Theta$.*

In the following, we assume the previous hypotheses are satisfied for $m = 2$ and we denote $\Omega_R^e = \Omega_0^e \cap B_R$.

ii) *For all $p \in (2, \bar{p}_1)$ where \bar{p}_1 is introduced in Remark 3.17, for all $\xi \in \Theta$, we have*

$$\|\hat{U}'_0\|_{H^{1,0}(\Omega_0^e)} \leq C_1 \|\xi\|_{W^{1,2p/(p-2)}(\Omega_R^e)}, \quad (69)$$

$$\left| \int_{\Omega_0^e} \nabla U_0 \cdot \hat{U}'_0 \right| \leq C_1 \|\xi\|_{W^{1,p/(p-2)}(\Omega_R^e)}, \quad (70)$$

$$\left| \int_{\Omega_0^e} \nabla U_0 \cdot \hat{U}''_0 \right| \leq C_2 \|\xi\|_{W^{1,2p/(p-2)}(\Omega_R^e)}^2, \quad (71)$$

where $\hat{U}'_0 = \hat{U}'_\theta(0)(\xi)$, $\hat{U}''_0 = \hat{U}''_\theta(0)(\xi, \xi)$, $C_i = C_i(\Omega_0, U_0, f, R, p)$.

Proof.

Step 1. The proof of i) is similar to the one of the same result in the interior, see i) in Theorem 3.9. For convenience we present the proof.

The proof is based on the implicit function theorem. After changing the variable $x = (I + \theta)(y)$, the weak form (68) in Ω_θ^e is transformed in Ω_0^e as follows

$$\int_{\Omega_0^e} \nabla \hat{U}_\theta \cdot M_\theta \cdot \nabla \varphi - \hat{f}_\theta \varphi = 0, \quad (72)$$

where $M_\theta = (T \cdot {}^t T) \det[Id + \nabla \theta]$, $T = [Id + \nabla \theta]^{-1}$ and $\hat{f} = f \circ (I + \xi(t)) \det[I + \nabla \theta]$. Consider F defined by

$$F : \begin{array}{l} \Theta \times H_0^{1,0}(\Omega_0^e) \\ (\theta, \hat{U}) \end{array} \mapsto \begin{array}{l} H^{-1,0}(\Omega_0^e) \\ \left[\varphi \mapsto \int_{\Omega_0^e} \nabla \hat{U} \cdot M_\theta \cdot \nabla \varphi - \hat{f}_\theta \varphi \right]. \end{array}$$

It is easy to check that F is well defined and of class C^m in a neighborhood of $(0, U_0) \in \Theta \times H_0^{1,0}(\Omega_0^e)$ (here we use the fact that f is with compact support), and

$$\partial_U F(0, U_0)(U) : \varphi \mapsto \int_{\Omega_0^e} (\nabla U \cdot \nabla \varphi) dx, \quad \forall U, \varphi \in H_0^{1,0}(\Omega_0^e).$$

Note that $\partial_U F(0, U_0)$ defines an isomorphism from $H_0^{1,0}(\Omega_0^e)$ to $H^{-1,0}(\Omega_0^e)$, see Proposition 3.16. Then, from implicit function Theorem, there exists a C^m map $\theta \mapsto U(\theta)$ such that $F(\theta, U(\theta)) = 0$ for $\|\theta\|_\Theta$ small. From the uniqueness of solution to (11) we obtain $\hat{U}_\theta = U(\theta)$, which proves the differentiability of $\theta \mapsto \hat{U}_\theta$.

Step 2. Again the proofs of (69)-(71) are similar to the proofs of (38)-(40) (except that the choice of suitable test functions is easier here). For the proof of (69) we differentiate (72) and get

$$\int_{\Omega_0^e} (\nabla \hat{U}'_\theta \cdot M_\theta \cdot \nabla \varphi) = \int_{\Omega_0^e} \hat{f}'_\theta \varphi - \nabla \hat{U}_\theta \cdot M'_\theta \cdot \nabla \varphi. \quad (73)$$

With $\theta = 0$ and $\varphi = \hat{U}'_0$, as $\text{Supp}(\xi) \subset B(0, R)$, from (73) we get

$$\begin{aligned} \|\nabla \hat{U}'_0\|_{L^2(\Omega_0^e)}^2 &\leq C_1 \int_{\Omega_R^e} (1 + |\nabla U_0|)(|\xi| + |\nabla \xi|)(|\hat{U}'_0| + |\nabla \hat{U}'_0|) \\ &\leq C_1(1 + \|U_0\|_{W^{1,p}(\Omega_R^e)})\|\xi\|_{W^{1,2p/(p-2)}(\Omega_R^e)}\|\nabla \hat{U}'_0\|_{L^2(\Omega_0^e)}, \end{aligned} \quad (74)$$

with $C_1 = C_1(\Omega_0, R, f, p)$, which proves (69).

Similarly, if we take $\theta = 0$ and $\varphi = U_0$ in (73) we get

$$\begin{aligned} \left| \int_{\Omega_0^e} \nabla \hat{U}'_0 \cdot \nabla U_0 \right| &\leq C_1 \int_{\Omega_R^e} (1 + |\nabla U_0|)(|\xi| + |\nabla \xi|)|\nabla U_0| \\ &\leq C_1(1 + \|U_0\|_{W^{1,p}(\Omega_R^e)})\|\xi\|_{W^{1,p/(p-2)}(\Omega_R^e)}, \end{aligned}$$

with $C_1 = C_1(\Omega_0, f, R, p)$, which proves (70).

Step 3. For the proof of (71) we differentiate (73) at $\theta = 0$ and take $\varphi = U_0$. It gives

$$\begin{aligned} \left| \int_{\Omega_0^e} \nabla \hat{U}''_0 \cdot \nabla U_0 \right| &\leq C_2 \int_{\Omega_R^e} (1 + |\nabla U_0|^2)(|\xi|^2 + |\nabla \xi|^2) + |\nabla U_0| |\nabla \xi| |\nabla \hat{U}'_0| \\ &\leq C_2 \left(\|U_0\|_{W^{1,p}(\Omega_R^e)} \|\nabla \xi\|_{L^{2p/(p-2)}(\Omega_R^e)} \|\nabla \hat{U}'_0\|_{L^2(\Omega_R^e)} + \right. \\ &\quad \left. (1 + \|U_0\|_{W^{1,p}(\Omega_R^e)}^2) \|\xi\|_{W^{1,2p/(p-2)}(\Omega_R^e)}^2 \right) \\ &\leq C_2(1 + \|U_0\|_{W^{1,p}(\Omega_R^e)}^2) \|\xi\|_{W^{1,2p/(p-2)}(\Omega_R^e)}^2, \end{aligned}$$

with $C_2 = C_2(\Omega_0, f, R, p)$, which completes the proof. \square

3.4.3 Estimate of shape derivatives of the energy

Theorem 3.19 *Let Ω_0 be as in (13). Assume it is also Lipschitz. Let $\theta \in \Theta$, $\Omega_0^e = \mathbb{R}^N \setminus \overline{\Omega_0}$, $\Omega_\theta^e = (I + \theta)(\Omega_0^e)$, $f \in W^{m,\infty}(\mathbb{R}^N)$, $m \in \mathbb{N}^*$, with $\text{supp}(f) \subset B_R$, $U_\theta \in H_0^{1,0}(\Omega_\theta^e)$ the solution of (68) and $\mathcal{E}(\theta) = \int_{\Omega_\theta} |\nabla U_\theta|^2$. Then \mathcal{E} is of class C^m near $\theta = 0 \in \Theta$ and for $m \geq 2$, for all $p \in (2, \bar{p}_1)$ with \bar{p}_1 defined in Remark 3.17, for all $\xi \in \Theta$, we have with $\Omega_R^e = \Omega_0^e \cap B_R$*

$$|\mathcal{E}'(0)(\xi)| \leq C_1 \|\xi\|_{W^{1,p/(p-2)}(\Omega_R^e)}, \quad (75)$$

$$|\mathcal{E}''(0)(\xi, \xi)| \leq C_2 \|\xi\|_{W^{1,2p/(p-2)}(\Omega_R^e)}^2, \quad (76)$$

where $C_i = C_i(\Omega_0, U_0, f, R, p)$.

Proof. The proof of the theorem is very similar to the one of Theorem 3.9. We present it briefly. Note that

$$\mathcal{E}(\theta) = \int_{\Omega_0^e} \nabla \hat{U}_\theta \cdot M_\theta \cdot \nabla \hat{U}_\theta. \quad (77)$$

From Theorem 3.18 it follows that \mathcal{E} is C^m in a neighborhood of $\theta = 0$.

Differentiating (77) at $\theta = 0$ in a direction $\xi \in \Theta$, and then using (70) gives

$$\begin{aligned} |\mathcal{E}'(0)(\xi)| &\leq C_1 \left(\left| \int_{\Omega_0^\varepsilon} \nabla U_0 \cdot \nabla \hat{U}'_0 \right| + \int_{\Omega_R^\varepsilon} |\nabla U_0|^2 |\nabla \xi| \right) \\ &\leq C_1 \|\xi\|_{W^{1,p/(p-2)}(\Omega_R^\varepsilon)}, \end{aligned} \quad (78)$$

with $C_1 = C_1(\Omega_0, U_0, R, p, f)$, which proves (75).

Differentiating twice (77) at $\theta = 0$ in the direction ξ , and then using (69) and (71) gives

$$\begin{aligned} |\mathcal{E}''(0)(\xi, \xi)| &\leq C_2 \left| \int_{\Omega_0^\varepsilon} \nabla \hat{U}''_0 \cdot \nabla U_0 \right| + \int_{\Omega_0^\varepsilon} |\nabla \hat{U}'_0|^2 + |\nabla U_0|^2 (|\xi|^2 + |\nabla \xi|^2) \\ &\leq C_2 \|\xi\|_{W^{1,2p/(p-2)}(\Omega_0^\varepsilon)}^2, \end{aligned} \quad (79)$$

with $C_2 = C_2(\Omega_0, U_0, R, p, f)$, which completes the proof. \square

4 Application to optimal convex shapes

In this section, we remind the strategy from [20], and emphasize an application of the second order shape derivatives estimates obtained in Section 2.2, first in dimension two, then in higher dimensions.

4.1 Application in the planar case

Here is the main result of this subsection:

Theorem 4.1 *Let Ω_0 be an open convex subset of \mathbb{R}^2 , solution of the optimization problem:*

$$\min \left\{ J(\Omega) = R(E(\Omega), |\Omega|) - P(\Omega), \quad \Omega \subset \mathbb{R}^2 \text{ open, convex and such that } \partial\Omega \subset \{x, a \leq |x| \leq b\} \right\}, \quad (80)$$

where $R : \mathbb{R}^2 \rightarrow \mathbb{R}$ is smooth, $(a, b) \in (0, \infty]^2$, $a < b$, and E is an energy like (5), corresponding to the interior problem, or like (10), corresponding to the exterior problem.

Then every connected component of $(\partial\Omega_0)_{in} := \partial\Omega_0 \cap \{x, a < |x| < b\}$ is a finite union of straight segments.

We insist here on the fact that the existence of an optimal shape for problem (80) is true and easy to obtain in the case $0 < a < b < \infty$, see Remark 4.7. The cases $b = \infty$ and/or $a = 0$ require more attention: it may happens that minimizing sequences are not bounded (for example if $R = 0$), or that they converge to a segment.

Remark 4.2 This result was obtained in [20] for the two following particular cases:

- $E = E_f$ is the Dirichlet energy associated to $f \in H_{loc}^2(\mathbb{R}^2)$:

$$E(\Omega) = \int_{\Omega} \left(\frac{1}{2} |\nabla U_{\Omega}|^2 - f U_{\Omega} \right), \quad \text{where } U_{\Omega} \text{ solves } \begin{cases} -\Delta U_{\Omega} = f \text{ in } \Omega \\ U_{\Omega} \in H_0^1(\Omega) \end{cases}$$

- $E = \lambda_1$ is the first Dirichlet eigenvalue of Ω :

$$\lambda_1(\Omega) = \min \left\{ \frac{\int_{\Omega} |\nabla U|^2}{\int_{\Omega} U^2}, \quad U \in H_0^1(\Omega) \setminus \{0\} \right\}.$$

Remark 4.3 In Theorem 4.1, we chose a simple function of $P(\Omega)$. Actually, the same result would extend to functionals of the form $R(E(\Omega), |\Omega|, \lambda_1(\Omega), P(\Omega))$ with $p \rightarrow R(\cdot, \cdot, \cdot, p)$ being decreasing and concave.

Remark 4.4 Similarly to [20, Theorem 4, Example 8], Theorem 4.1 is also valid with a volume constraint. More precisely, any solution of

$$\min\{J(\Omega) = R(E(\Omega)) - P(\Omega), \Omega \subset \mathbb{R}^2 \text{ open, convex and such that } |\Omega| = V_0\} \quad (81)$$

(where $R : \mathbb{R} \rightarrow \mathbb{R}$ is smooth and E is as in Theorem 4.1) is a polygon. See also Remark 4.6 for other geometrical constraints.

In order to analyze problem (80), we use, as in [19, 20], the following classical parametrization of 2-dimensional convex with polar coordinates $(r, \theta) \in [0, \infty) \times \mathbb{T}$, where $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$:

$$\Omega_u := \left\{ (r, \theta) \in [0, \infty) \times \mathbb{R} ; r < \frac{1}{u(\theta)} \right\}, \quad (82)$$

where u is a positive and 2π -periodic function. A simple computation shows that the curvature of Ω_u is

$$\kappa_{\partial\Omega_u} = \frac{u'' + u}{\left(1 + \left(\frac{u'}{u}\right)^2\right)^{3/2}}. \quad (83)$$

This implies that Ω_u is convex if and only if $u'' + u \geq 0$, which has to be understood in the sense of $H^{-1}(\mathbb{T})$ if u is not C^2 . More precisely, if $u \in H^1(\mathbb{T})$ then $u'' + u \geq 0$ if and only if

$$\forall v \in H^1(\mathbb{T}) \quad \text{with } v \geq 0, \quad \int_{\mathbb{T}} (uv - u'v') d\theta \geq 0.$$

Throughout this section, any function defined on \mathbb{T} is considered as the restriction to \mathbb{T} of a 2π -periodic function on \mathbb{R} , with the same regularity.

With this parametrization, considering $j(u) = J(\Omega_u)$, Problem (80) is equivalent to

$$\min \{j(u), u'' + u \geq 0, u \in \mathcal{U}_{ad}\}, \quad \text{where } \mathcal{U}_{ad} = \{u \in W^{1,\infty}(\mathbb{T}), 1/u \in [a, b]\}. \quad (84)$$

Then we have the following result proven in [20, Theorem 3] where $|v|_{H^1(\mathbb{T})}^2 = \int_{\mathbb{T}} (v')^2 d\theta$.

Theorem 4.5 *Let $u_0 > 0$ be a solution for (84) and $\mathbb{T}_{in} := \left\{ \theta \in \mathbb{T}, a < \frac{1}{u_0(\theta)} < b \right\}$. Assume $j : W^{1,\infty}(\mathbb{T}) \rightarrow \mathbb{R}$ is C^2 and that there exist $s \in [0, 1)$, $\alpha > 0$, $\beta, \gamma \in \mathbb{R}$ such that, for any $v \in W^{1,\infty}(\mathbb{T})$, we have*

$$j''(u_0)(v, v) \leq -\alpha |v|_{H^1(\mathbb{T})}^2 + \gamma |v|_{H^1(\mathbb{T})} \|v\|_{H^s(\mathbb{T})} + \beta \|v\|_{H^s(\mathbb{T})}^2. \quad (85)$$

If I is a connected component of \mathbb{T}_{in} , then

$$u_0'' + u_0 \text{ is a finite sum of Dirac masses in } I.$$

This result, combined with the estimates from Section 2.2 will lead to a proof of Theorem 4.1. Indeed, formula (83) explains that Ω_u is polygonal if and only if $u'' + u$ is a sum of Dirac masses.

Remark 4.6 As in [20], we may focus on a geometrical constraint different from $\partial\Omega \subset \{x, a \leq |x| \leq b\}$. In that case, the previous results remain valid when replacing the definitions of $\mathbb{T}_{in}, (\partial\Omega_0)_{in}$ with:

$$\mathbb{T}_{in} = \{\theta \in \mathbb{T} / \exists \varepsilon, \delta > 0, \forall v \in W_0^{1,\infty}(\theta - \varepsilon, \theta + \varepsilon) \text{ with } \|v\|_{W^{1,\infty}} < \delta, u_0 + v \in \mathcal{U}_{ad}\}.$$

$$(\partial\Omega_0)_{in} = \left\{ x \in \partial\Omega_0 / \exists \theta \in \mathbb{T}_{in}, x = \frac{1}{u_0(\theta)}(\cos \theta, \sin \theta) \right\}.$$

where \mathcal{U}_{ad} replaces $\{\Omega / \partial\Omega \subset \{x, a \leq |x| \leq b\}\}$ and includes the geometrical constraint, except the convexity one. Note that $(\partial\Omega_0)_{in}$ describes the part of the boundary which is *inside* the constraints, apart from the convexity one. However, if one deals also with a finite-dimensional equality constraint (for example $|\Omega| = V_0$), then one should work with a Lagrange multiplier, see Remark 4.4.

Remark 4.7 Due to the convexity constraint, the existence of a solution to (80) can be proved easily when $0 < a < b < \infty$. We briefly provide a proof here: let $(\Omega_n)_{n \in \mathbb{N}} \in \mathcal{O}_{ad}^{\mathbb{N}}$, with parametrization $u_n \in \mathcal{U}_{ad}$, be a minimizing sequence of $J(\Omega)$. From $u_n'' + u_n \geq 0$ and $1/u_n \in [a, b]$, it follows that $(u_n)_{n \in \mathbb{N}}$, is strongly relatively compact in $H^1(\mathbb{T})$ and therefore we may assume $\lim_{n \rightarrow \infty} u_n = u_0$ in $H^1(\mathbb{T})$ (see also Remark 4.12), with $u_0'' + u_0 \geq 0$ and $1/u_0 \in [a, b]$. In such conditions the domains Ω_n satisfy a uniform exterior cone condition and will converge to Ω_0 for the Hausdorff convergence, with parametrization u_0 . Then U_{Ω_n} will converge to U_{Ω_0} in $H^1(B(0, b))$ (see [12, Prop 2.4.4 and Theorem 3.2.13]), and thanks to the continuity of the perimeter for the $H^1(\mathbb{T})$ -norm we get that $P(\Omega_n)$ converges to $P(\Omega_0)$, which implies that Ω_0 solves (80).

Proof of Theorem 4.1. Let u_0 such that $\Omega_0 = \Omega_{u_0}$. Then u_0 is a solution of (84), where $j(u) = R(e(u), m(u)) - p(u)$ and $e(u) = E(\Omega_u), m(u) = |\Omega_u|, p(u) = P(\Omega_u)$.

The second order derivatives related to the geometrical terms $m(u)$ and $p(u)$ can be easily computed. Indeed, first note that for every $u \in W^{1,\infty}(\mathbb{T}), u > 0$ we have

$$m(u) = \int_{\mathbb{T}} \frac{1}{2u^2} d\theta, \quad p(u) = \int_{\mathbb{T}} \frac{\sqrt{u^2 + u'^2}}{u^2} d\theta.$$

It follows that m and p are twice differentiable in $W^{1,\infty}(\mathbb{T})$ and there exist some real numbers β_1, β_2, γ (depending on u_0) and $\alpha > 0$ such that,

$$\begin{cases} |m''(u_0)(v, v)| \leq \beta_1 \|v\|_{L^2(\mathbb{T})}^2 \\ p''(u_0)(v, v) \geq \alpha |v|_{H^1(\mathbb{T})}^2 - \gamma |v|_{H^1(\mathbb{T})} \|v\|_{L^2(\mathbb{T})} - \beta_2 \|v\|_{L^2(\mathbb{T})}^2, |v|_{H^1(\mathbb{T})}^2 = \int_{\mathbb{T}} (v')^2 d\theta. \end{cases}$$

It remains the difficult term $e''(u_0)(v, v)$, which relies on Section 2.2. To handle this term, we consider $v \in W^{1,\infty}(\mathbb{T})$ and introduce the vector field:

$$\begin{aligned} \xi &\in C^2((-t_0, t_0), W^{1,\infty}(\mathbb{R}^2, \mathbb{R}^2)), \\ \xi(t) &= \left(\frac{1}{u+tv} - \frac{1}{u} \right) e^{i\theta} \eta(r, \theta) \quad \text{on } \partial\Omega_0 \text{ (in polar coordinates),} \end{aligned} \tag{86}$$

where $\eta \in C_c^\infty(\mathbb{R}^2) = \mathcal{D}(\mathbb{R}^2), 0 \leq \eta \leq 1, \eta = 1$ (resp. $\eta = 0$) in a neighborhood of $\partial\Omega_0$ (resp. the origin). Let us point out that $(I + \xi(t))(\Omega_u) = \Omega_{u+tv}, e(u+tv) = E(\Omega_{u+tv}) = \mathcal{E}(\xi(t))$ and also that

$$\forall v \in W^{1,\infty}(\mathbb{T}) : \quad \xi'(0)(v) = -\frac{v}{u^2} e^{i\theta}, \quad \xi''(0)(v, v) = 2\frac{v^2}{u^3} e^{i\theta} \quad \text{on } \partial\Omega. \tag{87}$$

Then, we will differentiate twice around 0 the map $t \mapsto \varphi(t) := E(\Omega_{u+tv}) = \mathcal{E}(\xi(t))$:

$$\begin{aligned} \varphi'(0) &= e'(u_0)(v) = \mathcal{E}'(0)(\xi'(0)), \\ \varphi''(0) &= e''(u_0)(v, v) = \mathcal{E}''(0)(\xi'(0), \xi'(0)) + \mathcal{E}'(0)(\xi''(0)). \end{aligned}$$

and therefore Corollary 2.6 (see also (22), (23)) implies there exists $\varepsilon \in (0, 1/2)$ such that

$$|e''(u_0)(v, v)| \leq \beta_3(\varepsilon) \|\xi''(0)\|_{H^{\frac{1}{2}-\varepsilon}(\partial\Omega)} + \beta_4(\varepsilon) \|\xi'(0)\|_{H^{1-\varepsilon}(\partial\Omega)}^2.$$

We then use the fact that the $H^{1-\varepsilon}(\partial\Omega)$ and $H^{1-\varepsilon}(\mathbb{T})$ norms are equivalent (since the transformation $\psi = \psi(r, \theta) := \frac{r}{u(\theta)} e^{i\theta}$ is bi-Lipschitz near \mathbb{T} and $\psi(\mathbb{T}) = \partial\Omega$), and the fact that $H^{1-\varepsilon}(\mathbb{T})$ is a Banach algebra to obtain

$$|e''(u)(v, v)| \leq \beta_5(\varepsilon) \|v\|_{H^{1-\varepsilon}(\mathbb{T})}^2,$$

Easy computations about the term $R(E(\Omega), |\Omega|)$ imply that j'' satisfies (85), and therefore Theorem 4.5 applies. It follows that $u_0'' + u_0$ is a finite sum of Dirac masses in any connected component of \mathbb{T}_{in} . Geometrically speaking, considering the formula (83), this correspond to the fact that any connected component of $(\partial\Omega_0)_{in}$ is polygonal and concludes the proof. \square

4.2 Application in the multi-dimensional case

In the multi-dimensional case, convexity constraint in shape optimization is much less understood, see [3, 11] and the work of T. Lachand-Robert. Nevertheless, we can use a parametrization similar to the one used in Section 4.1, and we show in this section that our estimates of shape derivatives allows to obtain results in any dimension.

For $N \geq 2$, if $u : \mathbb{S}^{N-1} \rightarrow (0, \infty)$ is given, $\mathbb{S}^{N-1} = \{x \in \mathbb{R}^N, |x| = 1\}$, we can consider

$$\Omega_u := \left\{ (r, \theta) \in [0, \infty) \times \mathbb{S}^{N-1}, \quad r < \frac{1}{u(\theta)} \right\}. \quad (88)$$

The function u is the so-called gauge function of Ω_u . The set Ω_u is convex if and only if the 1-homogeneous extension of u , denoted by the same letter and given by $u(x) = |x|u(x/|x|)$ is convex in \mathbb{R}^N (in this section, we will refer to this property by saying that $u : \mathbb{S}^{N-1} \rightarrow \mathbb{R}$ is convex), see [28, Section 1.7] for example. In this way, we describe every bounded convex open set containing the origin. Throughout this section, the regularity of any function defined on \mathbb{S}^{N-1} is seen as the regularity on $\mathbb{R}^N \setminus \{0\}$ of its 1-homogeneous extension, and it is classical that it is equivalent to the regularity of the set Ω_u itself.

With this parametrization, considering $j(u) = J(\Omega_u)$, problem (1) is equivalent to

$$\min \{ j(u), u : \mathbb{S}^{N-1} \rightarrow (0, \infty) \text{ convex}, u \in \mathcal{U}_{ad} \}, \quad (89)$$

where $\mathcal{U}_{ad} \subset W^{1,\infty}(\mathbb{S}^{N-1})$ is defined as $\mathcal{U}_{ad} = \{u \in W^{1,\infty}(\mathbb{S}^{N-1}), \Omega_u \in \mathcal{O}_{ad}\}$ and \mathcal{O}_{ad} is the class of admissible open bounded sets considered in (1) (not taking into account the convexity constraint).

Then in the same spirit as Theorem 4.5, we can prove the following where we denote $|v|_{H^1(\mathbb{S}^{N-1})}^2 = \int_{\mathbb{S}^{N-1}} |\nabla_\tau v|^2 d\theta$, $\nabla_\tau =$ tangential gradient on \mathbb{S}^{N-1} :

Theorem 4.8 *Let $u_0 > 0$ be a solution for (89), where \mathcal{U}_{ad} is a convex subset of $W^{1,\infty}(\mathbb{S}^{N-1})$. Assume $j : W^{1,\infty}(\mathbb{S}^{N-1}) \rightarrow \mathbb{R}$ is C^2 and that there exists $s \in [0, 1)$, $\alpha > 0$, $\beta, \gamma \in \mathbb{R}$ such that for any $v \in W^{1,\infty}(\mathbb{S}^{N-1})$ we have*

$$j''(u_0)(v, v) \leq -\alpha |v|_{H^1(\mathbb{S}^{N-1})}^2 + \gamma |v|_{H^1(\mathbb{S}^{N-1})} \|v\|_{H^s(\mathbb{S}^{N-1})} + \beta \|v\|_{H^s(\mathbb{S}^{N-1})}^2. \quad (90)$$

Then the set

$$T_{u_0} = \{v \in W^{1,\infty}(\mathbb{S}^{N-1}) / \exists \varepsilon > 0, \forall |t| < \varepsilon, u_0 + tv \in \mathcal{U}_{ad} \text{ and is convex}\}, \quad (91)$$

is a linear vector space of finite dimension.

Proof. It is easy to check that T_{u_0} is a linear vector space. Since u_0 is optimal, we have $j''(u_0)(v, v) \geq 0$ for every $v \in T_{u_0}$. To conclude, we prove that the unit ball of T_{u_0} for the $H^s(\mathbb{S}^{N-1})$ -norm is relatively compact. Indeed, thanks to (90), we get

$$\forall v \in T_{u_0} \text{ such that } \|v\|_{H^s(\mathbb{S}^{N-1})} \leq 1, \quad \alpha|v|_{H^1(\mathbb{S}^{N-1})}^2 \leq \gamma|v|_{H^1(\mathbb{S}^{N-1})} + \beta,$$

which easily leads to

$$\forall v \in T_{u_0} \text{ such that } \|v\|_{H^s(\mathbb{S}^{N-1})} \leq 1, \quad |v|_{H^1(\mathbb{S}^{N-1})} \leq \frac{|\gamma| + \sqrt{\gamma^2 + 4\alpha|\beta|}}{2\alpha}.$$

From the compact embedding of $H^s(\mathbb{S}^{N-1})$ in $H^1(\mathbb{S}^{N-1})$, we conclude that the unit ball of $T_{u_0} \subset H^s(\mathbb{S}^{N-1})$ has a compact closure and therefore from Riesz Theorem, T_{u_0} has finite dimension. \square

As a corollary, the results we obtained in this paper, namely iii), Theorem 2.3, lead to the following generalization and improvement of [3, Theorem 4.5]:

Corollary 4.9 *Let Ω_0 be an open convex subset of \mathbb{R}^N , solution of the optimization problem:*

$$\min \left\{ J(\Omega) = R(E(\Omega), |\Omega|) - P(\Omega), \quad \Omega \subset \mathbb{R}^N \text{ open, convex and such that } \partial\Omega \subset \{x, a \leq |x| \leq b\} \right\}, \quad (92)$$

where $R : \mathbb{R}^2 \rightarrow \mathbb{R}$ is smooth, $(a, b) \in (0, \infty]^2$, $a < b$, and E is an energy like (5), corresponding to the interior problem.

Then denoting by u_0 the gauge function of Ω_0 , T_{u_0} defined in (91) has finite dimension. In particular, if ω is a C^2 relatively open subset of $(\partial\Omega_0)_{in} := \partial\Omega_0 \cap \{x, a < |x| < b\}$, then the Gauss curvature of Ω_0 vanishes on ω .

Compared to [3, Theorem 4.5], we have enlarged the class of functionals under consideration, and we require less regularity on the optimal set.

Remark 4.10 Contrary to Theorem 4.1, we cannot deal with the exterior problem when $N > 2$. This is due to the fact that for a Lipschitz domain Ω in dimension $N \geq 3$, our estimates do not imply that there exists $s < 1$ such that $E''(\Omega)(\xi, \xi) \leq \|\xi\|_{H^s(\partial\Omega)}$ (whereas it is the case for $N = 2$, see (23)).

Remark 4.11 As in Remark 4.6, we may focus on a geometrical constraint different from $\partial\Omega \subset \{x, a \leq |x| \leq b\}$. In that case, given \mathcal{O}_{ad} a class of admissible open bounded set, the previous result remain valid when replacing the definition of $(\partial\Omega_0)_{in}$ as follows

$$\begin{aligned} (\partial\Omega_0)_{in} &= \left\{ x \in \partial\Omega_0 / \exists \theta \in \mathbb{S}_{in}^{N-1}, x = \frac{1}{u_0(\theta)}\theta \right\}, \text{ with} \\ \mathbb{S}_{in}^{N-1} &= \left\{ \theta \in \mathbb{S}^{N-1} / \exists \varepsilon, \delta > 0, \forall v \in W_0^{1,\infty}(B_{\mathbb{S}^{N-1}}(\theta, \varepsilon)) \text{ with } \|v\|_{W^{1,\infty}} < \delta, u_0 + v \in \mathcal{U}_{ad} \right\} \text{ and} \\ \mathcal{U}_{ad} &= \{u, \Omega_u \in \mathcal{O}_{ad}\}, \end{aligned}$$

if \mathcal{U}_{ad} is convex, as assumed in Theorem 4.8.

Remark 4.12 As in Remark 4.7, the existence of a solution to (92) can be proved easily when $0 < a < b < \infty$. The proof is similar to the one in Remark 4.7, the only adaptation to the multi-dimensional case is the fact that a sequence of 1-homogeneous functions $u_n : \overline{B_1} \rightarrow \mathbb{R}$ (where B_1 is the unit ball of \mathbb{R}^N) such that $\forall n \in \mathbb{N}, \forall \theta \in \mathbb{S}^{N-1}, \alpha \leq u_n(\theta) \leq \beta$, with $0 < \alpha < \beta < \infty$, is strongly relatively compact in $H^1(\overline{B_1})$ (actually in $W^{1,p}(\overline{B_1})$ for any $p < \infty$). To that end, we first notice that $\|\nabla u_n\|_\infty \leq \beta$ (see for example [5]), which

implies that up to a subsequence u_n converges to u for the $L^\infty(\overline{B_1})$ -norm. Clearly u is then 1-homogeneous, convex, and satisfies $\forall \theta \in \mathbb{S}^{N-1}, \alpha \leq u(\theta) \leq \beta$. To obtain the convergence for the H^1 -norm, we follow [2, Lemma 2.1.2]. As in Remark 4.7, the perimeter, the volume, and the energy E are continuous, and therefore the set whose gauge function is the limit of u_n is a minimizer.

Proof of Corollary 4.9. Let u_0 such that $\Omega_0 = \Omega_{u_0}$. Then u_0 is a solution of (89), where $j(u) = R(e(u), m(u)) - p(u)$ and $e(u) = E(\Omega_u), m(u) = |\Omega_u|, p(u) = P(\Omega_u)$.

The second order derivatives related to the geometrical terms $m(u)$ and $p(u)$ can be easily computed. Indeed, first note that for every $u \in W^{1,\infty}(\mathbb{S}^{N-1}), u > 0$ we have (see for instance [7])

$$m(u) = \int_{\mathbb{S}^{N-1}} \frac{1}{Nu^N} d\theta, \quad p(u) = \int_{\mathbb{S}^{N-1}} \frac{\sqrt{u^2 + |\nabla_\tau u|^2}}{u^N} d\theta.$$

where ∇_τ denotes the tangential gradient on \mathbb{S}^{N-1} . It follows that m and p are twice differentiable and there exists β_1 (depending on u_0) such that,

$$|m''(u_0)(v, v)| \leq \beta_1 \|v\|_{L^2(\mathbb{T})}^2$$

Easy direct calculations give

$$\begin{aligned} p'(u)(v) &= \int_{\mathbb{S}^{N-1}} \partial_u G(u, |\nabla_\tau u|) v + \partial_p G(u, |\nabla_\tau u|) \frac{\nabla_\tau u \cdot \nabla_\tau v}{|\nabla_\tau u|}, \\ p''(u)(v, v) &= \int_{\mathbb{S}^{N-1}} \partial_{uu} G(u, |\nabla_\tau u|) v^2 + 2\partial_{up} G(u, |\nabla_\tau u|) \frac{\nabla_\tau u \cdot \nabla_\tau v}{|\nabla_\tau u|} v + \\ &\quad \partial_{pp} G(u, |\nabla_\tau u|) \frac{(\nabla_\tau v \cdot \nabla_\tau u)^2}{|\nabla_\tau u|^2} + \partial_p G(u, |\nabla_\tau u|) \frac{|\nabla_\tau u|^2 |\nabla_\tau v|^2 - (\nabla_\tau v \cdot \nabla_\tau u)^2}{|\nabla_\tau u|^3}. \end{aligned}$$

Taking into account $\partial_p G(u, p) = \frac{p}{u^N \sqrt{u^2 + p^2}}, \partial_{pp} G(u, p) = \frac{u^2}{u^N (u^2 + p^2)^{3/2}}$ and rearranging the last two terms gives

$$\begin{aligned} p''(u)(v, v) &= \int_{\mathbb{S}^{N-1}} \partial_{uu} G(u, |\nabla_\tau u|) v^2 + 2\partial_{up} G(u, |\nabla_\tau u|) \frac{\nabla_\tau u \cdot \nabla_\tau v}{|\nabla_\tau u|} v + \\ &\quad \frac{1}{u^N \sqrt{u^2 + |\nabla_\tau u|^2}} \left[|\nabla_\tau v|^2 - \frac{|\nabla_\tau u|^2}{u^2 + |\nabla_\tau u|^2} \left(\frac{\nabla_\tau u}{|\nabla_\tau u|} \cdot \nabla_\tau v \right)^2 \right]. \end{aligned}$$

As the first and second terms are controlled by $\beta_2 \|v\|_{L^2}^2$ and $\gamma \|v\|_{L^2} \|v\|_{H^1}$ respectively, with β_2, γ depending on u_0 , and

$$|\nabla_\tau v|^2 - \frac{|\nabla_\tau u|^2}{u^2 + |\nabla_\tau u|^2} \left(\frac{\nabla_\tau u}{|\nabla_\tau u|} \cdot \nabla_\tau v \right)^2 \geq (1 - (1 - \alpha)) |\nabla_\tau v|^2 = \alpha |\nabla_\tau v|^2,$$

with $\alpha = \alpha(u_0) > 0$, it follows

$$p''(u_0)(v, v) \geq \alpha \|v\|_{H^1(\mathbb{S}^{N-1})}^2 - \gamma \|v\|_{H^1(\mathbb{S}^{N-1})} \|v\|_{L^2(\mathbb{S}^{N-1})} - \beta_2 \|v\|_{L^2(\mathbb{S}^{N-1})}^2.$$

As in the proof of Theorem 4.1, the term $e''(u_0)(v, v)$ relies on Section 2.2, this time in any dimension, but restricted to the case of convex domains. With the same proof and using iii), Theorem 2.3, we obtain

$$|e''(u_0)(v, v)| \leq \beta_3(r) \|\xi''(0)\|_{W^{1-\frac{1}{r}, r}(\partial\Omega)} + \beta_4(r) \|\xi'(0)\|_{W^{1-\frac{1}{2r}, 2r}(\partial\Omega)}^2,$$

valid for any $r \in (1, \infty)$. We would like to write the above inequality in terms of the $H^s(\mathbb{S}^{N-1})$ -norm of v , for some $s \in (0, 1)$. For this, first we note that any bi-Lipschitz transformation from \mathbb{S}^{N-1} to $\partial\Omega_0$ (in our case it is $\theta \mapsto (1/u(\theta), \theta)$) defines a diffeomorphism from $W^{s,p}(\partial\Omega_0)$ to $W^{s,p}(\mathbb{S}^{N-1})$, for every $s \in [0, 1]$ and $p \in [1, \infty]$. Then the above inequality is equivalent to

$$|e''(u_0)(v, v)| \leq \beta_5(r, u) \left\| \frac{v^2}{u_0^3} \right\|_{W^{1-\frac{1}{r}, r}(\mathbb{S}^{N-1})} + \beta_6(r, u) \left\| \frac{v}{u_0^2} \right\|_{W^{1-\frac{1}{2r}, 2r}(\mathbb{S}^{N-1})}^2, \quad (93)$$

Note that here we used the formulas (87) with $e^{i\theta}$ replaced by θ (of norm 1).

For the first term in (93) we wonder for which $s \in (0, 1)$ the following inclusion is continuous:

$$H^s(\mathbb{R}^{N-1})H^s(\mathbb{R}^{N-1})W^{1,\infty}(\mathbb{R}^{N-1}) \subset W^{1-1/r, r}(\mathbb{R}^{N-1}). \quad (94)$$

We use [27, Corollary, p. 189], which refers to the more general embedding $B_{p_1, q_1}^{s_1} \cdots B_{p_m, q_m}^{s_m} \subset B_{p, q}^s$ in \mathbb{R}^N . Note that by using a partition of unity, some Lipschitz transformations and the extension theorem, we may use these embeddings as if \mathbb{S}^{N-1} was \mathbb{R}^{N-1} , and therefore we use it with $N-1$ instead of N . In our specific case, this leads to the condition

$$\left(1 - \frac{1}{r}\right) \frac{N}{2} < s < 1. \quad (95)$$

Clearly, for every $s \in (0, 1)$ one can find r so that (95) holds, which implies that (94) is valid.

For the second term in (93), we look for the inclusion

$$H^s(\mathbb{R}^{N-1})W^{1,\infty}(\mathbb{R}^{N-1}) \subset W^{1-1/(2r), 2r}(\mathbb{R}^{N-1}). \quad (96)$$

which is valid (again using [27]) if

$$1 - \frac{1}{2r} + \frac{N-1}{2} \left(1 - \frac{1}{r}\right) < s < 1. \quad (97)$$

Inequality (97) has a solution r (sufficiently close to 1) for any $s \in (1/2, 1)$, which implies that (96) holds for every $s \in (1/2, 1)$.

Combining (94) and (96) with (93) gives

$$|e''(u)(v, v)| \leq \beta_7(s, u_0) \|v\|_{H^s(\mathbb{S}^{N-1})}^2,$$

for any $s \in (1/2, 1)$.

Easy computations about the term $R(E(\Omega), |\Omega|)$ implies then that $j''(u_0)$ satisfies (90), and therefore Theorem 4.8 applies, and T_{u_0} has finite dimension.

The fact that the Gauss curvature must vanish where it is defined is an easy consequence. Indeed, if ω is as in the statement of Corollary 4.9 and if the Gauss curvature is positive at some point of ω , then it is positive in a neighborhood $\hat{\omega}$ of this point. As a consequence, any smooth function with compact support in $\hat{\omega}$ is in T_{u_0} . This contradicts the fact that T_{u_0} has finite dimension. \square

References

- [1] F. ANDREU, J. M. MAZÓN, J. D. ROSSI, *The best constant for the Sobolev trace embedding from $W^{1,1}(\Omega)$ into $L^1(\Omega)$* , *Nonlinear Analysis* 59 (2004) 1125-1145
- [2] D. BUCUR, G. BUTTAZZO, *Variational Methods in Shape Optimization Problems*, Birkhäuser, 2005

- [3] D. BUCUR, I. FRAGALÀ, J. LAMBOLEY, *Optimal convex shapes for concave functional*, ESAIM Control Optim. Calc. Var. 18 (2012), no. 3, 693–711
- [4] H. BREZIS, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Springer 2011.
- [5] A. FIGALLI AND F. MAGGI, *On the shape of liquid drops and crystals in the small mass regime*. *Arch. Rat. Mech. Anal.*, 201:143–207, 2011.
- [6] S. J. FROMM, *Potential space estimates for Green potentials in convex domains*, Proc. of AMS, Vol. 119, No. 1 (1993)
- [7] B. FUGLEDE, *Stability in the isoperimetric problem for convex or nearly spherical domains in \mathbb{R}^n* , Trans. of the AMS, Vol 314, Number 2, 1989
- [8] D. GILBARG, N. TRUDINGER, *Elliptic partial differential equations of second order*, Springer, 1998.
- [9] A. B. GULISASHVILI, *Traces of functions belonging to Sobolev and Besov spaces and extensions from subsets of the Euclidean space*, Journal of Soviet Mathematics, March 1989, Volume 44, Issue 6, pp 843-849]
- [10] P. GRISVARD, *Elliptic problems in nonsmooth domains*, Pitman, Boston, 1985.
- [11] E. HARRELL, A. HENROT, J. LAMBOLEY, *On the local minimizers of the Mahler volume*, Journal of convex analysis, 2015.
- [12] A. HENROT, M. PIERRE, *Variation et optimisation de formes : une analyse géométrique*, Springer 2005.
- [13] D. JERISON, C. KENIG, *The Inhomogeneous Dirichlet Problem in Lipschitz Domains*, Journal of Functional Analysis, vol. 130, no. 1 (1995), 161-219, 1995.
- [14] H. JIA, D. LI, L. WANG, *Global regularity for divergence form elliptic equations on quasiconvex domains*, J. Differential Equations 249 (2010), no. 12, 3132–3147.
- [15] A. JONSSON, H. WALLIN, *Function Spaces on Subsets of R^n* , Harwood Acad. Publ., London (1984) Mathematical Reports, Volume 2, Part 1.
- [16] J. KADLEC, *The regularity of the solution of the Poisson problem in a domain whose boundary is similar to that of a convex domain*, Czechoslovak Math. J. 89 (1964), pp. 386-393.
- [17] S. LANG, *Real And Functional Analysis*, Graduate texts in Mathematics, Third Edition, Springer.
- [18] O. LADYZHENSKAYA, N. URAL'TSEVA, *Linear and quasilinear elliptic equations*, Academic Press, 1968.
- [19] J. LAMBOLEY, A. NOVRUZI, *Polygons as optimal shapes under convexity constraint*, SIAM J. Control Optim. 48 (2009/10), no. 5, 30033025
- [20] J. LAMBOLEY, A. NOVRUZI, M. PIERRE, *Regularity and Singularities of Optimal Convex Shapes in the Plane*, *Archive for rational mechanics and analysis*, Volume 205, Issue 1 (2012), 311-343.
- [21] J. MARSCHALL, *The trace of Sobolev-Slobodeckij spaces on Lipschitz domains*, Manuscripta Mathematica, vol. 58, no. 1 (1987), 47-65.
- [22] D. MITREA, M. MITREA, *General Second Order, Strongly Elliptic Systems in Low Dimensional Non-smooth Manifolds*, Contemporary Math., Vol. 277 (2001), 61-86.

- [23] D. MITREA, M. MITREA, L. YAN, *Boundary value problems for the Laplacian in convex and semiconvex domains*, JFA, 258 (2010) 2507-2585
- [24] M. MITREA, M. TAYLOR, *Potential theory on Lipschitz domains in Riemannian manifolds: Sobolev-Besov space results and the Poisson problem*, J. Funct. Anal., 176:1 (2000), 1979.
- [25] M. MITREA, M. TAYLOR, *Boundary Layer Methods for Lipschitz Domains in Riemannian Manifolds*, Journal of Functional Analysis 163, 181-251 (1999)
- [26] J. C. NÉDÉLEC, *Approximation des équations intégrales en physiques et mécanique*, Centres de Mathématiques Appliquées, Ecole Polytechnique, 1977.
- [27] TH. RUNST, W. SICKEL, *Sobolev spaces of fractional order, Nemytsckij operators, and nonlinear Partial Differential Equations*, Walter de Gruyter, Berlin, New York, 1996
- [28] R. SCHNEIDER, *Convex bodies: the Brunn-Minkowski Theory* Encyclopedia of Mathematics and its Applications, 44. Cambridge University Press, Cambridge, 1993
- [29] J. SIMON, *Differentiation with respect to the domain in boundary value problems*, Numer. Funct. Anal. Optim., 2, no 7-8 (1980), 649-687.
- [30] ZH. SHEN, *Bounds of Riesz Transforms on L^p Spaces for Second Order Elliptic Operators*, Annales de l'Institut Fourier, Tome 55, no 1 (2005), p. 173-197.
- [31] H. TRIEBEL, *Theory of function spaces, III*, Monographs in Mathematics, Birkhäuser, 2006.