

**CAHIER DU LAMSADE**

**172**

Juin 2000

Differential approximation results for traveling  
salesman problem

Jérôme Monnot, Vangelis Th. Paschos, Sophie Toulouse

## Table of Contents

Résumé	ii
Abstract	ii
<b>1 Introduction</b>	<b>1</b>
<b>2 Preserving differential approximation for several min_TSP versions</b>	<b>3</b>
<b>3 2_OPT and differential approximation for the general minimum traveling salesman</b>	<b>4</b>
<b>4 Approximating min_TSP12</b>	<b>7</b>
4.1 Specification of the min_TSP12-algorithm and evaluation of $\lambda(K_n)$ . . . . .	8
4.1.1 Construction and evaluation of $C$ . . . . .	9
4.1.1.1 $q = 1$ . . . . .	10
4.1.1.2 $q = k$ . . . . .	10
4.1.1.3 $q = k + 1$ . . . . .	10
4.1.2 Construction and evaluation of $D$ . . . . .	10
4.1.3 Construction and evaluation of $\tilde{T}$ . . . . .	11
4.1.4 Overall specification of the min_TSP12-algorithm, construction and evaluation of $T$ . . . . .	11
4.2 A bound for $\omega(K_n)$ . . . . .	11
4.2.1 Disjoint elementary paths on $V(C)$ . . . . .	12
4.2.2 Disjoint elementary paths on $V(D)$ . . . . .	12
4.2.3 Disjoint elementary paths on $V(\hat{T})$ . . . . .	12
4.2.3.1 $q > 0$ . . . . .	12
4.2.3.2 $q = 0$ . . . . .	13
4.2.4 Disjoint elementary paths on $V(K_n)$ . . . . .	13
4.2.4.1 $q > 0,  E_2  <  C_{p+1} $ . . . . .	13
4.2.4.2 $q > 0,  E_2  =  C_{p+1} $ . . . . .	13
4.2.4.3 $q = 0,  E_2  <  C_{p+1} $ . . . . .	14
4.3 The differential approximation ratio of TSP12 . . . . .	15
4.4 Ratio 3/4 is tight for TSP12 . . . . .	15
<b>5 Further results for minimum traveling salesman</b>	<b>15</b>
5.1 Bridges between differential and standard approximation . . . . .	15
5.2 An inapproximability result . . . . .	17
<b>6 Differential approximation of maximum traveling salesman</b>	<b>18</b>
<b>7 An improvement of the standard ratio for the maximum traveling salesman with distances 1 and 2</b>	<b>19</b>
<b>8 Towards stronger differential-inapproximability results for the traveling salesman</b>	<b>19</b>

# Résultats d'approximation différentielle pour le problème du voyageur du commerce

## Résumé

Nous commençons par démontrer que les versions maximisation et minimisation du problème du voyageur de commerce sont approximables à rapport différentiel  $1/2$ . Nous présentons ensuite une  $3/4$ -approximation polynomiale du cas particulier à distances 1 et 2; ce résultat nous permet notamment de ramener le rapport standard connu pour la version maximisation de ce sous-problème de  $5/7$  à  $7/8$ . Nous proposons enfin un résultat négatif: approximer le voyageur de commerce, à coût minimum comme maximum, à mieux que  $3475/3476 + \epsilon$  est **NP**-difficile pour tout  $\epsilon > 0$ .

**Mots-clé** : algorithme d'approximation, rapport d'approximation, problème **NP**-complet, complexité, réduction, voyageur du commerce.

## Differential approximation results for traveling salesman problem

### Abstract

We prove that both minimum and maximum traveling salesman problems can be approximately solved, in polynomial time within approximation ratio bounded above by  $1/2$ . We next prove that, when dealing with edge-distances 1 and 2, both versions are approximable within  $3/4$ . Based upon this result, we then improve the standard approximation ratio known for maximum traveling salesman with distances 1 and 2 from  $5/7$  to  $7/8$ . Finally, we prove that, for any  $\epsilon > 0$ , it is **NP**-hard to approximate both problems within better than  $3475/3476 + \epsilon$ .

**Keywords**: approximation algorithm, approximation ratio, **NP**-complete problem, complexity, reduction, traveling salesman.

## 1 Introduction

Given a complete graph on  $n$  vertices, denoted by  $K_n$ , with positive distances on its edges, the minimum traveling salesman problem (min\_TSP) consists in minimizing the cost of a Hamiltonian cycle, the cost of such a cycle being the sum of the distances on its edges. The maximum traveling salesman problem (max\_TSP) consists in maximizing the cost of a Hamiltonian cycle. Further special but very natural cases of TSP are the ones where edge-distances are defined using the  $\ell_2$  norm (Euclidean TSP), or where edge-distances verify triangle inequalities (metric TSP); an interesting sub-case of the metric TSP is the one in which edge-distances are only 1 or 2 (TSP12). Both min\_ and max\_TSP, even in their restricted versions mentioned just mentioned above, are famous NP-hard problems.

In general, NP optimization (NPO) problems are commonly defined as follows.

**Definition 1.** An NPO problem  $\Pi$  is as a four-tuple  $(\mathcal{I}, \mathbf{S}, v_I, \text{opt})$  such that:

1.  $\mathcal{I}$  is the set of instances of  $\Pi$  and it can be recognized in polynomial time;
2. given  $I \in \mathcal{I}$  (let  $|I|$  be the size of  $I$ ),  $\mathbf{S}(I)$  denotes the set of feasible solutions of  $I$ ; moreover, there exists a polynomial  $P$  such that, for every  $S \in \mathbf{S}(I)$  (let  $|S|$  be the size of  $S$ ),  $|S| = O(P(|I|))$ ; furthermore, given any  $I$  and any  $S$  with  $|S| = O(P(|I|))$ , one can decide in polynomial time if  $S \in \mathbf{S}(I)$ ;
3. given  $I \in \mathcal{I}$  and  $S \in \mathbf{S}(I)$ ,  $v_I(S)$  denotes the value of  $S$ ;  $v_I$  is integer, polynomially computable and is commonly called objective function;
4.  $\text{opt} \in \{\max, \min\}$ . ■

Given an instance  $I$  of an NPO problem  $\Pi$  and a polynomial time approximation algorithm  $\mathbf{A}$  feasibly solving  $\Pi$ , we will denote by  $\omega(I)$ ,  $\lambda_{\mathbf{A}}(I)$  and  $\beta(I)$  the values of the worst solution of  $I$ , of the approximated one (provided by  $\mathbf{A}$  when running on  $I$ ), and the optimal one for  $I$ , respectively. There exist mainly two thought processes dealing with polynomial approximation. Commonly ([13]), the quality of an approximation algorithm for an NP-hard minimization (resp., maximization) problem  $\Pi$  is expressed by the ratio (called standard in what follows)  $\rho_{\mathbf{A}}(I) = \lambda(I)/\beta(I)$ , and the quantity  $\rho_{\mathbf{A}} = \inf\{r : \rho_{\mathbf{A}}(I) < r, I \text{ instance of } \Pi\}$  (resp.,  $\rho_{\mathbf{A}} = \sup\{r : \rho_{\mathbf{A}}(I) > r, I \text{ instance of } \Pi\}$ ) constitutes the approximation ratio of  $\mathbf{A}$  for  $\Pi$ . Recent works ([9, 8]), strongly inspired by [3] (see also [12, 23]), bring to the fore another approximation measure, as powerful as the traditional one (concerning the type, the diversity and the quantity of the results produced), the ratio (called differential in what follows)  $\delta_{\mathbf{A}}(I) = (\omega(I) - \lambda(I))/(\omega(I) - \beta(I))$ . The quantity  $\delta_{\mathbf{A}} = \sup\{r : \delta_{\mathbf{A}}(I) > r, I \text{ instance of } \Pi\}$  is the differential approximation ratio of  $\mathbf{A}$  for  $\Pi$ . In what follows, we use notation  $\rho$  when dealing with standard ratio and notation  $\delta$  when dealing with the differential one. Moreover  $\rho(\Pi)$  (resp.,  $\delta(\Pi)$ ) will denote the best standard (resp., differential) approximation ratio for  $\Pi$ .

In [3], the term “trivial solution” is used to denote what in [9, 8] and here is called *worst solution*. Moreover, all the examples in [3] carry over NP-hard problems for which worst solution can be trivially computed. This is for example the case of maximum independent set where, given a graph, the worst solution is the empty set, or of minimum vertex cover, where the worst solution is the vertex-set of the input-graph, or even of the minimum graph-coloring where one can trivially color the vertices of the input-graph using a distinct color per vertex. On the contrary, for TSP things are very different. Let us take for example min\_TSP. Here, given a graph  $K_n$ , the worst solution for  $K_n$  is a maximum total-distance Hamiltonian cycle, i.e., the optimal solution of max\_TSP in  $K_n$ . The computation of such a solution is very far from being trivial since max\_TSP is NP-hard. Obviously, the same holds when one considers max\_TSP

and tries to compute a worst solution for its instance. In order to remove ambiguities about the concept of the worst solution, the following definition, proposed in [9], will be used here.

**Definition 2.** Given a typical instance  $I$  of an **NPO** problem  $\Pi$ , the worst solution of  $I$  is the optimal solution of a new **NPO** problem  $\bar{\Pi}$  where items 1 to 3 of definition 1 are identical for both  $\Pi$  and  $\bar{\Pi}$ , and

$$\text{opt}(\bar{\Pi}) = \begin{cases} \max & \text{opt}(\Pi) = \min \\ \min & \text{opt}(\Pi) = \max \quad \blacksquare \end{cases}$$

One of the features of the differential ratio is to be stable under affine transformation of the objective function of a problem and so it does not create a dissymmetry between minimization and maximization problems. This is very clear in the case of TSP. Dealing with  $\text{min\_TSP}$  it is very well-known that its general version is not approximable in polynomial time within better than  $2^{p(n)}$  for a polynomial  $p$ . On the other hand, its maximization version,  $\text{max\_TSP}$ , the **NP**-hardness of which is immediately proved if one replaces distance  $d(i, j)$  for  $\text{min\_TSP}$  by  $M - d(i, j)$  in  $\text{max\_TSP}$ , for an  $M$  greater than the largest edge distance in the input graph of  $\text{min\_TSP}$ , can be approximated in polynomial time within  $5/7$  ([20]).

Let us recall some standard terminology from the theory of the polynomial approximation of the **NP**-hard problems (for the standard approximation framework). Given an **NP** minimization (resp., maximization) problem  $\Pi$ , a *constant-ratio approximation algorithm* for  $\Pi$  is a polynomial time approximation algorithm (PTAA) guaranteeing approximation ratio bounded above (resp., below) by a fixed constant, i.e., by a constant that does not depend on any input-parameter of  $\Pi$ . **APX** is the class of the **NP** optimization problems solved by constant-ratio PTAA's. A *polynomial time approximation schema* (PTAS) for  $\Pi$  is a sequence of PTAA's (receiving as inputs any instance of  $\Pi$  and a fixed constant  $\epsilon$ ) guaranteeing approximation ratio bounded above (resp., below) by  $1 + \epsilon$  (resp.,  $1 - \epsilon$ ), for every  $\epsilon > 0$ . If a PTAS is polynomial in both  $n$  and  $1/\epsilon$ , then it is called *fully polynomial time approximation schema* (FPTAS). For the differential approximation, the ratio achieved by polynomial time approximation schemata is  $1 - \epsilon$  for both minimization and maximization. Finally, **APX**-complete is the class of problems in **APX**, which, in addition, are complete with respect to the existence of a PTAS solving them, in other words, if any **APX**-complete problem could be solved by a PTAS, then any other **APX**-complete problem could be so.

As it is shown in [9, 8], many problems behave in completely different ways regarding traditional or differential approximation. This is, for example, the case of minimum graph-coloring or, even, of minimum vertex-covering. This paper deals with another example of the diversity in the nature of approximation results achieved within the two frameworks, the TSP. For this problem and its versions mentioned above, a bunch of standard-approximation results (positive or negative) have been obtained until nowadays. The first inapproximability result is the one of [21] (see also [13]) affirming that it is **NP**-hard to approximate  $\text{min\_TSP}$  within any constant factor; with the same proof, one can easily refine the result of [21] to deduce the inapproximability of  $\text{min\_TSP}$  within any ratio of the form  $2^{p(n)}$  for any polynomial  $p$ . On the other hand, the metric  $\text{min\_TSP}$  is approximable within  $3/2$  ([5]), the symmetric  $\text{min\_TSP12}$  within  $7/6$  ([18]) (recall that the original proof of the **NP**-completeness of the  $\text{min\_TSP}$  is done by reduction to  $\text{min\_TSP12}$ ), while the asymmetric version of  $\text{min\_TSP12}$  is approximable within  $17/12$  ([22]). Moreover,  $\text{min\_TSP12}$  is **APX**-complete ([18]), consequently, given the result of [2], it cannot be solved by a PTAS unless **P=NP**; in other words,  $\exists \epsilon > 0$  for which approximation of  $\text{min\_TSP12}$  within ratio smaller than  $1 + \epsilon$  is **NP**-hard. Furthermore, even in graphs where the density of the subgraph spanned by the edges of length 1 is bounded below by a constant  $c \in ]0, 1/2[$ ,  $\text{min\_TSP12}$  cannot be solved by a polynomial time approximation schema ([11]). The works of [10] and more recently of [4] refine the result of [18] specifying

for  $\epsilon$ . In [4] is proved that for any  $\epsilon > 0$ , it is **NP**-hard to approximate  $\text{min\_TSP12}$  within ratio smaller than, or equal to,  $3475/3476 - \epsilon$ ; in other words the result of [10] gives a value – equal to  $1/3476 - \epsilon'$ ,  $\forall \epsilon' > 0$  – for the hardness threshold  $\epsilon$  of  $\text{min\_TSP12}$  refining so the negative results of [18, 10]. Finally another restrictive version of the metric  $\text{min\_TSP}$ , the Euclidean  $\text{min\_TSP}$  can be solved by a standard PTAS ([1]). A complete list of standard-approximation results for  $\text{min\_TSP}$  is given in [6].

In what follows, we show that, in the differential approximation framework the classical  $2\_OPT$  algorithm, originally devised in [7] and revisited in numerous works (see, for example, [15]), approximately solves  $\text{min\_TSP}$  with edge-distances bounded by a polynomial of  $n$  within differential approximation ratio  $1/2$ . In other words,  $2\_OPT$  provides for these graphs solutions “fairly close” to the optimal and, simultaneously, “fairly far” from the worst one. We also prove that, in the opposite of what happens in the standard framework, metric  $\text{min\_TSP}$  and general  $\text{min\_TSP}$  are equi-approximable in the differential framework. Moreover we prove that  $\text{min\_TSP12}$  is approximable within  $3/4$ .

For  $\text{max\_TSP}$  things are much more optimistic for standard approximation, since this problem is in **APX**. By the end of 70s it has been proved in [12] that  $2\_OPT$  guarantees approximation ratio  $1/2$  for  $\text{max\_TSP}$ . More recently, in [20] is proved that  $\text{max\_TSP}$  can be solved by a standard PTAA within ratio  $5/7$ , if the distance-vector is symmetric and within  $38/63$ , if it is asymmetric. The dissymmetry in the approximability of  $\text{min\_}$  and  $\text{max\_TSP}$  can be considered as somewhat curious given the structural symmetry existing between them. In fact the transformation  $d \mapsto M - d$  mentioned above and revisited in detail in section 6 is affine. Since differential approximation is stable under affine transformation of the objective function,  $\text{min\_TSP}$  and  $\text{max\_TSP}$  are equi-approximable.

In what follows, we will denote by  $V = \{v_1, \dots, v_n\}$  the vertex-set of  $K_n$ , by  $E$  its edge-set and, for  $v_i v_j \in E$ , we denote by  $d(v_i, v_j)$  (or by  $d(i, j)$  when no ambiguity occurs) the distance of the edge  $v_i v_j \in E$ ; we consider that the distance-vector is symmetric and integer. Given a feasible TSP-solution  $T(K_n)$  of  $K_n$  (both  $\text{min\_}$  and  $\text{max\_TSP}$  have the same set of feasible solutions), we denote by  $d(T(K_n))$  its (objective) value;  $T$  will be indexed by  $\text{min}$  or  $\text{max}$  depending on whether it deals with  $\text{min\_}$  or  $\text{max\_TSP}$ . When necessary, the values of the worst case solution, the approximated one and the optimal one for  $\text{min\_TSP}$  ( $\text{max\_TSP}$ ) will be denoted by  $\omega_{\text{min}}(K_n)$ ,  $\lambda_{\mathbf{A}}^{\text{min}}(K_n)$  and  $\beta_{\text{min}}(K_n)$  ( $\omega_{\text{max}}(K_n)$ ,  $\lambda_{\mathbf{A}}^{\text{max}}(K_n)$  and  $\beta_{\text{max}}(K_n)$ ), respectively. Given a graph  $G$  induced by  $K_n$ , we denote by  $V(G)$  its vertex-set. Finally, given any set  $C$  of edges, we denote by  $d(C)$  the total distance of  $C$ , i.e., the quantity  $\sum_{v_i v_j \in C} d(i, j)$ .

## 2 Preserving differential approximation for several $\text{min\_TSP}$ versions

This section is a preliminary one containing several results about how differential approximation is preserved between several restrictive versions of  $\text{min\_TSP}$ .

**Proposition 1.** *Metric  $\text{min\_TSP}$  and general  $\text{min\_TSP}$  are differentially equi-approximable.*

**Proof.** Obviously, metric  $\text{min\_TSP}$  being a special case of the general one, can be solved within the same differential approximation ratio with the latter.

Suppose now that metric  $\text{min\_TSP}$  is approximately solved within differential ratio  $\delta$ . Given an instance  $I = (K_n, \vec{d})$  of general  $\text{min\_TSP}$  (set  $d_{\text{max}} = \max\{d(i, j) : v_i v_j \in E\}$ ), one can transform it into a new one  $I' = (K_n, \vec{d}')$  by changing,  $\forall v_i v_j \in E$ , distance  $d(i, j)$  of the former to  $d'(i, j) = d_{\text{max}} + d(i, j)$ . It is easy to see that  $I'$  is metric, and that every feasible tour  $T(I)$  of  $(K_n, \vec{d})$  remains feasible for  $(K_n, \vec{d}')$ . The cost of such a tour becomes  $d(T(I')) = d(T(I)) + nd_{\text{max}}$ . Then the  $\delta$ -PTAA for metric  $\text{min\_TSP}$  will achieve

$$\delta = \frac{\omega(I') - d(T(I'))}{\omega(I') - \beta(I')} = \frac{\omega(I) + nd_{\text{max}} - d(T(I)) - nd_{\text{max}}}{\omega(I) + nd_{\text{max}} - \beta(I) - nd_{\text{max}}} = \frac{\omega(I) - d(T(I))}{\omega(I) - \beta(I)}.$$

Consequently, every PTAA for metric  $\text{min\_TSP}$  can simultaneously solve general  $\text{min\_TSP}$  within the same differential approximation ratio. ■

Let  $d_{\min} = \min\{d(i, j) : v_i v_j \in E\}$ . Then, if one transforms every distance  $d(i, j)$  into  $d(i, j) - d_{\min} + 1$ , one obtains a complete graph where  $d_{\min} = 1$  and with arguments completely analogous to the ones of proposition 1, the following holds.

**Proposition 2.** *General  $\text{min\_TSP}$  and  $\text{min\_TSP}$  with  $d_{\min} = 1$  are differentially equi-approximable.*

We next consider another class of instances, the one where the edge-distances are either  $a$ , or  $b$  (notorious member of this class of  $\text{min\_TSP}$ -problems, denoted by  $\text{min\_TSP}ab$ , is the  $\text{min\_TSP}12$ ). Suppose, without loss of generality that  $a < b$ . Then, by proposition 2,  $\text{min\_TSP}ab$  is equi-approximable with  $\text{min\_TSP}1b$ . Consider now an instance of the latter problem. If one sets  $b = 2$  for all the  $b$ -edges (edges of distance  $b$ ), then by arguments completely similar to the ones of the proof of proposition 1 (and since for a tour  $T$  containing  $k_b$   $b$ -edges,  $d(T) = n + (b - 1)k_b$ ), the following result holds.

**Proposition 3.**  *$\text{min\_TSP}ab$  and  $\text{min\_TSP}12$  are differentially equi-approximable.*

Note that results analogous to the ones of propositions 1, 2 and 3 do not hold in the standard approximation framework.

### 3 2\_OPT and differential approximation for the general minimum traveling salesman

In what follows, we denote by **D-APX** the analogous of the class **APX**, the class of **NPO** problems solved by a constant-ratio PTAA, for the differential approximation framework.

**Theorem 1.**  *$\text{min\_TSP}$  is differentially approximable within approximation ratio  $1/2$  and this ratio is tight.*

**Proof.** In what follows, suppose that a tour is listed as the set of its edges and consider the following algorithm of [7].

```
BEGIN /2_OPT/
(1)  start from any feasible tour T;
(2)  REPEAT
(3)      pick a new set  $\{v_i v_j, v_{i'} v_{j'}\} \subset T$ ;
(4)      IF  $d(i, j) + d(i', j') > d(i, i') + d(j, j')$  THEN  $T \leftarrow (T \setminus \{v_i v_j, v_{i'} v_{j'}\}) \cup \{v_i v_{i'}, v_j v_{j'}\}$  FI
(5)  UNTIL no improvement of  $d(T)$  is possible;
(6)  OUTPUT T;
END. /2_OPT/
```

Suppose now that, starting from a vertex denoted by  $v_1$ , the rest of the vertices is ordered following the tour  $T$  finally computed by 2\_OPT (so, given a vertex  $v_i$ ,  $i = 1, \dots, n - 1$ ,  $v_{i+1}$  is its successor with respect to  $T$ ;  $v_{n+1} = v_1$ ). Let us fix one optimal tour and denote it by  $T^*$ . Given a vertex  $v_i$ , denote by  $v_{s^*(i)}$  its successor in  $T^*$  (remark that  $v_{s^*(i)+1}$  is the successor of  $v_{s^*(i)}$  in  $T$ ; in other words, edge  $v_{s^*(i)} v_{s^*(i)+1} \in T$ ). Finally let us fix one (of the eventually many) worst-case (maximum total-distance) tour  $T_\omega$ .

The tour  $T$  computed by 2\_OPT is a local optimum for the 2-exchange of edges in the sense that every interchange between two non-intersecting edges of  $T$  and two non-intersecting edges of  $E \setminus T$  will produce a tour of total distance at least equal to  $d(T)$ . This implies in particular that,  $\forall i \in \{1, \dots, n\}$ ,

$$d(i, i + 1) + d(s^*(i), s^*(i) + 1) \leq d(i, s^*(i)) + d(i + 1, s^*(i) + 1);$$

so, writing the expression above for all  $i \in \{1, \dots, n\}$ , we get

$$\sum_{i=1}^n (d(i, i+1) + d(s^*(i), s^*(i)+1)) \leq \sum_{i=1}^n (d(i, s^*(i)) + d(i+1, s^*(i)+1)) \quad (1)$$

Moreover, it is easy to see that the following holds:

$$\bigcup_{i=1, \dots, n} \{v_i v_{i+1}\} = \bigcup_{i=1, \dots, n} \{v_{s^*(i)} v_{s^*(i)+1}\} = T \quad (2)$$

$$\bigcup_{i=1, \dots, n} \{v_i v_{s^*(i)}\} = T^* \quad (3)$$

$$\bigcup_{i=1, \dots, n} \{v_{i+1} v_{s^*(i)+1}\} = \text{some feasible tour } T' \quad (4)$$

Let us show that  $T' = \bigcup_{i=1, \dots, n} \{v_{i+1} v_{s^*(i)+1}\}$  is feasible. Recall that an acyclic permutation is a bijective function  $f : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$  such that,  $\forall i \in \{1, \dots, n\}$ :

$$\begin{cases} f^{(k)}(i) \neq i & k < n \\ f^{(n)}(i) = i \end{cases}$$

Every feasible tour  $T$ , oriented as mentioned above, can be seen as an acyclic permutation. Consider now the following mappings

$$\begin{aligned} s^* & : i \mapsto s^*(i) \\ f & : i \mapsto i+1 \\ h & : i \mapsto s^*(i-1)+1. \end{aligned}$$

It is easy to see that if  $s^*$  is an acyclic permutation and  $f$  is a permutation, then  $h = f \circ s^* \circ f^{-1}$  is an acyclic permutation. Moreover, it is not hard to see that pairs  $(i, h(i))$  correspond  $(\text{mod } n)$  to the edges of  $T'$ .

Combining expression (1) with expressions (2), (3) and (4), one gets:

$$\begin{aligned} (2) & \implies \sum_{i=1}^n d(i, i+1) + \sum_{i=1}^n d(s^*(i), s^*(i)+1) = 2\lambda_{2\_OPT}(K_n) \\ (3) & \implies \sum_{i=1}^n d(i, s^*(i)) = \beta(K_n) \\ (4) & \implies \sum_{i=1}^n d(i+1, s^*(i)+1) = d(T') \leq \omega(K_n) \end{aligned} \quad (5)$$

and expressions (1) and (5) lead to

$$2\lambda_{2\_OPT}(K_n) \leq \beta(K_n) + \omega(K_n) \iff \frac{\omega(K_n) - \lambda_{2\_OPT}(K_n)}{\omega(K_n) - \beta(K_n)} \geq \frac{1}{2}.$$

Consequently,  $\delta_{2\_OPT} \geq 1/2$ .

Consider now a  $K_{4n+8}$ ,  $n \geq 0$ , set  $V = \{v_i : i = 1, \dots, 4n+8\}$ , let

$$\begin{aligned} d(2k+1, 2k+2) &= 1 & k &= 0, 1, \dots, 2n+3 \\ d(4k+2, 4k+4) &= 1 & k &= 0, 1, \dots, n+1 \\ d(4k+3, 4k+5) &= 1 & k &= 0, 1, \dots, n \\ d(4n+7, 1) &= 1 \end{aligned}$$



and set the distances of all the remaining edges to 2. Then,

$$\begin{aligned}
T &= \{v_i v_{i+1} : i = 1, \dots, 4n + 7\} \cup \{v_{4n+8} v_1\} \\
T^* &= \{v_{2k+1} v_{2k+2} : k = 0, \dots, 2n + 3\} \cup \{v_{4k+2} v_{4k+4} : k = 0, \dots, n + 1\} \\
&\quad \cup \{v_{4k+3} v_{4k+5} : k = 0, \dots, n\} \cup \{v_{4n+7} v_1\} \\
T_w &= \{v_{2k+2} v_{2k+3} : k = 0, \dots, 2n + 2\} \cup \{v_{2k+1} v_{2k+4} : k = 0, \dots, 2n + 1\} \\
&\quad \cup \{v_{2k+2} v_{2k+5} : k = 0, \dots, 2n + 1\} \cup \{v_{4n+8} v_1\}.
\end{aligned}$$

In figure 1,  $T^*$  and  $T_w$  are shown for  $n = 1$  ( $T = \{1, \dots, 11, 12, 1\}$ ). Hence,  $\delta_{2\_OPT}(K_{4n+8}) = 1/2$  and this completes the proof of the theorem. ■




Figure 1. Tightness of the 2\\_OPT approximation ratio for  $n = 1$ .

From the proof of the tightness of the ratio of 2\\_OPT, the following corollary is immediately deduced.

**Corollary 1.**  $\delta_{2\_OPT} = 1/2$  is tight even for  $\min\_TSP_{12}$ .

A first case of polynomial complexity for algorithm 2\\_OPT (even if edge-distances of the graph are exponential in  $n$ ) is for graphs where the number of (feasible) tour-values, denoted by  $\sigma(K_n)$ , is polynomial in  $n$ . Here, since there exists a polynomial number of different  $\min\_TSP$  solution-values, achievement of a locally minimal solution (starting, at worst for the worst-value solution) will need a polynomial number of steps (at most  $\sigma(K_n)$ ) for 2\\_OPT.

Theorem 1 obviously works in polynomial time when  $d_{\max}$  is bounded above by a polynomial of  $n$ . However, even when this condition is not satisfied, there exist restrictive cases of  $\min\_TSP$  for which 2\\_OPT remains polynomial.

Consider now complete graphs with a fixed number  $k \in \mathbb{N}$  of distinct edge-distances,  $d_1, d_2, \dots, d_k$ . Then, any tour-value can be seen as  $k$ -tuple  $(n_1, n_2, \dots, n_k)$  with  $n_1 + n_2 + \dots + n_k = n$ , where  $n_1$  edges of the tour are of distance  $d_1, \dots, n_k$  edges are of distance  $d_k$  ( $\sum_{i=1}^k n_i d_i = d(T)$ ). Consequently, the consecutive solutions retained by 2\\_OPT (in line (4)) before attaining a local minimum are, at most, as many as the number of the arrangements with repetitions of  $k$  distinct items between  $n$  items (in other words, the number of all the distinct  $k$ -tuples formed by all the numbers in  $\{1, \dots, n\}$ ), i.e., bounded above by  $O(n^k)$ .

Another class of polynomially solved instances is the one where  $\beta(K_n) = O(P(n))$  where  $P$  is a polynomial of  $n$ . Recall that, from proposition 1, general and metric  $\min\_TSP$  are differentially equi-approximable. Consequently, given an instance  $K_n$  where  $\beta(K_n)$  is polynomial,  $K_n$  can be transformed into a graph  $K'_n$  as in proposition 1. Then, if one runs the algorithm of [5] in order to obtain an initial feasible tour  $T$  (line (1) of algorithm  $2\_OPT$ ), then its total distance, at most  $3/2$  times the optimal one, will be of polynomial value and, consequently,  $2\_OPT$  will need a polynomial number of steps until attaining a local minimum.

Let us note that the first and the fourth of the above cases cannot be decided in polynomial time. However, if one systematically transforms general  $\min\_TSP$  into a metric one (proposition 1) and then uses the algorithm of [5] in line (1) of  $2\_OPT$ , then all instances meeting the second item of corollary 2 will be solved in polynomial time even if we cannot recognize them.

**Corollary 2.** *The following versions of  $\min\_TSP$  are in  $D-APX$  (solved by  $2\_OPT$  within ratio  $1/2$ ):*

- on graphs where the optimal tour-value is polynomial in  $n$ ;
- on graphs where the number of feasible tour-values is polynomial in  $n$  (examples of these graphs are the ones where edge-distances are polynomially bounded, or even the ones where there exists a fixed number of distinct edge-distances).

## 4 Approximating $\min\_TSP12$

Let us first recall that, given a graph  $G$ , a 2-matching is a set  $M$  of edges of  $G$  such that if  $V(M)$  is the set of the endpoints of  $M$ , the vertices of the graph  $(V(M), M)$  have degree at most 2, in other words, the graph  $(V(M), M)$  is a collection of cycles and simple paths. A 2-matching is optimal if it is the largest over all the 2-matchings of  $G$ . As it is shown in [14], *an optimal triangle-free 2-matching can be computed in polynomial time.*

Our  $\min\_TSP12$  PTAA is based upon a special kind of triangle-free 2-matching in  $K_n$ , the cycles of which will be progressively patched in order to produce a Hamiltonian tour. In what follows, we deal with optimal triangle-free 2-matchings, i.e., with *triangle-free collections of cycles*.

**Theorem 2.**  *$\min\_TSP12$  is approximable within differential approximation ratio  $\delta \geq 3/4$ . This ratio is tight for the algorithm devised.*

**Proof.** Let  $M = (C_1, C_2, \dots)$  be any maximal triangle-free 2-matching of  $K_n$ . In the sequel, we call by *value of a 2-matching* the sum of the distances of its edges. For any matching  $M$ , we will denote its value by  $d(M)$ . Also, let us call *cycle-patching* (see also [18]) the operation consisting in taking two cycles  $C_i$  and  $C_j$  of  $M$ , in picking edges  $v_k v_l \in C_i$ ,  $v_p v_q \in C_j$  and in transforming  $C_i$ , and  $C_j$  into a unique cycle  $C = C_i \cap C_j \setminus \{v_k v_l, v_p v_q\} \cup \{e_{ij}, e'_{ij}\}$ , where  $\{e_{ij}, e'_{ij}\} = \{v_k v_p, v_l v_q\}$ , or  $\{e_{ij}, e'_{ij}\} = \{v_k v_q, v_l v_p\}$ . This specifies the following procedure, polynomial in  $n$ , computing, in addition, the total distance of the cycle resulting from cycle patching.

```
BEGIN /CYCLE_PATCH/
  take edges  $v_k v_l \in C_i$  and  $v_p v_q \in C_j$ ;
   $C_{ij}^1 \leftarrow C_i \cup C_j \setminus \{v_k v_l, v_p v_q\} \cup \{v_k v_p, v_l v_q\}$ ;
   $C_{ij}^2 \leftarrow C_i \cup C_j \setminus \{v_k v_l, v_p v_q\} \cup \{v_k v_q, v_l v_p\}$ ;
  OUTPUT  $C_{ij} \leftarrow \operatorname{argmin}\{d(C_{ij}^1), d(C_{ij}^2)\}$ ;
END. /CYCLE_PATCH/
```

#### 4.1 Specification of the `min_TSP12`-algorithm and evaluation of $\lambda(K_n)$

In the sequel, we will first specify a PTAA `min_TSP12` and estimate the value  $\lambda_{\text{TSP12}}(K_n) = d(T(K_n))$  of the Hamiltonian tour computed. Next, we will compute a lower bound for  $\omega(K_n)$ . As for theorem 1, we will exhibit a feasible tour of a certain value. Since worst solution's value is larger than the value of every other Hamiltonian tour of  $K_n$ , the value of the tour exhibited will be the bound claimed.

Let  $\hat{M}$  be an optimal triangle-free 2-matching of  $K_n$  (recall that, as we have mentioned, such a matching is maximal, i.e, it does only contains cycles). Starting from  $\hat{M}$ , one can easily construct an optimal 2-matching  $M^*$  where every patching of two cycles strictly increases its value. In what follows, we will call  $M^*$  *2-minimal*. Construction of  $M^*$  can be done in polynomial time by the following procedure.

```

BEGIN /2_MIN/
  Mp ← ∅;
  REPEAT
    pick a new set {Ci, Cj} ⊆  $\hat{M}$ ;
    FOR all vkvl ∈ Ci, vpvq ∈ Cj DO
      Mp ←  $\hat{M} \setminus \{C_i, C_j\} \cup \text{CYCLE\_PATCH}(C_i, C_j)$ 
      IF d( $\hat{M}$ ) > d(Mp) THEN  $\hat{M} \leftarrow M_p$  FI
    OD
  UNTIL no improvement of d( $\hat{M}$ ) is possible;
  OUTPUT M* ←  $\hat{M}$ ;
END. /2_MIN/

```

**Remark 1.** In any 2-minimal matching  $M$  there exists *at most* one cycle  $C$  containing 2-edges (edges of distance 2). In fact, if not, procedure `CYCLE_PATCHING` can always be applied in order to patch two distinct cycles containing 2-edges into one cycle with total distance no longer than the sum of the distances of the two cycles patched. Moreover, if  $M = C$ , then  $M$  is an optimal solution for `min_TSP` (in general, a Hamiltonian cycle being a particular triangle-free 2-matching,  $d(M) \leq \beta_{\min}(K_n)$ ). ■

Fix a 2-minimal triangle-free matching  $M^* = (C_1, C_2, \dots, C_{p+1})$  (recall that  $M^*$  is a minimum total-distance triangle-free 2-matching) and suppose, without loss of generality, that  $p > 0$  and that cycles  $C_1, \dots, C_p$  contain only 1-edges (edges of distance 1) and that only cycle  $C_{p+1}$  contains, eventually, some 2-edges. Finally, recall that it is assumed that  $|C_i| \geq 4$ . The following facts can be concluded regarding  $M^*$ .

**Fact 1.**  $\forall (C, C') \in M^* \times M^*$  such that  $C \neq C'$ ,  $\forall uv \in C$ ,  $\forall u'v' \in C'$ ,  $\max\{d(u, u'), d(v, v')\} = \max\{d(u, v'), d(v, u')\} = 2$ . ■

**Fact 2.** If vertex  $u$  is adjacent to a 2-edge in  $C_{p+1}$ , then,  $\forall u' \notin V(C_{p+1})$ ,  $d(u, u') = 2$ . ■

**Fact 3.** If  $uu'$  and  $vv'$  are two distinct non-adjacent 2-edges of  $C_{p+1}$ , then  $d(u, v) = d(u, v') = d(u'v) = d(u'v') = 2$ . ■

Given  $M^* = (C_1, \dots, C_{p+1})$ , we first perform the following preprocessing on  $C_1, \dots, C_p$ .

```

BEGIN /PREPROCESS/
  PR ← ∅;
  WHILE possible DO
    arbitrarily pick Ci, Cj ∈ M* \ {Cp+1} linked by at least one 1-edge;

```

```

    PR ← PR ∪ {Ci, Cj};
    M* ← M* \ {Ci, Cj};
  OD
  OUTPUT PR;
END. /PREPROCESS/

```

Suppose the WHILE loop of PREPROCESS executed  $q$  times and denote by  $\{C_1^s, C_2^s\}$  the cycles considered during the  $s$ th execution of the loop,  $s = 1, \dots, q$ . Then  $PR = \cup_{s=1}^q \{C_1^s, C_2^s\}$ . Set  $r = p - 2q$  and denote by  $D_t$ ,  $t = 1, \dots, r$ , the cycles in  $\{C_1, \dots, C_p\} \setminus \cup_{s=1}^q \{C_1^s, C_2^s\}$ . Under all this,

$$M^* = \left\{ \bigcup_{s=1}^q \{C_1^s, C_2^s\} \right\} \cup \left\{ \bigcup_{t=1}^r D_t \right\} \cup C_{p+1}.$$

The following facts hold and complete the above discussion.

**Fact 4.**  $2q + r \geq 1$ ; if  $2q + r = 1$  then  $C_{p+1} \neq \emptyset$ . ■

**Fact 5.**  $\forall s \in \{1, \dots, q\}, \forall \ell \in \{1, 2\}, \forall e \in C_\ell^s, d(e) = 1$ . ■

**Fact 6.**  $\forall t \in \{1, \dots, r\}, \forall e \in D_t, d(e) = 1$ . ■

**Fact 7.**  $\forall s \in \{1, \dots, q\}, \exists i^s \in V(C_1^s), \exists I^s \in V(C_2^s)$  such that  $d(i^s, I^s) = 1$ . ■

**Fact 8.**  $\forall (t, t') \in \{1, \dots, r\} \times \{1, \dots, r\}, t \neq t', \forall (u, v) \in V(D_t) \times V(D_{t'}), d(u, v) = 2$ . ■

In the sequel, for  $s = 1, \dots, q$ , we denote by  $a^s$  and  $b^s$  (resp.,  $A^s$  and  $B^s$ ) the vertices adjacent to  $i^s$  (resp.,  $I^s$ ) in  $C_1^s$  (resp.,  $C_2^s$ ). We set  $c = \sum_{s=1}^q (|C_1^s| + |C_2^s|)$ ,  $d = \sum_{t=1}^r |D_t|$ ,  $E2 = \{e \in C_{p+1} : d(e) = 2\}$ . Following these notations,  $n = c + d + |C_{p+1}|$  and, denoting by  $|E2|$  cardinality of the set  $E2$ ,

$$d(M^*) = n + |E2| \tag{6}$$

We are well-prepared now to describe the algorithm proposed. Informally, it first patches cycles  $C_1^s$  and  $C_2^s$  into a single cycle  $C^s$ ,  $s = 1, \dots, q$ . Next, it patches cycle  $C^1$  with  $C^2$  to produce a cycle  $C$  which will be patched with  $C^3$ , and so on, finally producing a single cycle  $C$ . It does so for the cycles  $D_t$ ,  $t = 1, \dots, r$ , producing a single cycle  $D$ . Then it patches  $C$  and  $D$  in order to produce a partial tour  $\tilde{T}$  and finally it patches  $\tilde{T}$  and  $C_{p+1}$  obtaining so the final TSP-tour  $T(K_n)$ .

#### 4.1.1 Construction and evaluation of $C$

Construction of  $C$  is performed by means of the following procedure.

```

BEGIN /C/
  FOR s ← 1 to q DO using edge  $i^s I^s$   $C^s \leftarrow \text{CYCLE\_PATCH}(C_1^s, C_2^s)$ ; OD
  C ← C1;
  FOR s ← 1 TO q − 1 DO
    replacing as many 2-edges as possible C ← CYCLE_PATCH(C, Cs+1);
  OD
  OUTPUT C;
END. /C/

```

The call of algorithm CYCLE\_PATCH in the first FOR-loop of C is a very slightly different variant of the corresponding procedure presented above where one imposes to the 1-edge  $i^s I^s$  (fact 7) to be one of the cross-edges entering cycle  $C^s$ .

**Lemma 1.** *The 2-matching  $(C^1, C^2, \dots, C^q)$  produced during the  $q$  executions of the first FOR-loop of algorithm  $\mathcal{C}$  has value  $d(C^1, C^2, \dots, C^q) = c + q$ .*

**Proof of lemma 1.** Patching of  $C_1^s$  and  $C_2^s$  into  $C^s$  is done using 1-edge  $i^s I^s$  (fact 7),  $s = 1, \dots, q$ . Consequently, only one 2-edge has been included in  $C^s$  (the one used with  $i^s I^s$  to patch  $C_1^s$  and  $C_2^s$ ). Such an edge always exists because of fact 1. So, for  $s = 1, \dots, q$ , execution of  $\text{CYCLE\_PATCH}(C_1^s, C_2^s)$  in the first FOR-loop of  $\mathcal{C}$  will produce in all exactly  $q$  2-edges replacing and  $q$  1-edges replacing  $2q$  1-edges. Consequently,  $d(C^1, C^2, \dots, C^q) = \sum_{s=1}^q (|C_1^s| + |C_2^s|) + q = c + q$  and this completes the proof of lemma 1. ■

During the executions of  $\text{CYCLE\_PATCH}$  in the second FOR-loop of  $\mathcal{C}$ , we try that the total distance of the resulting cycle is no longer than the sum of the total distances of the cycles patched. In other words, we try to not produce additional 2-edges in the resulting cycle. Here the following lemma holds.

**Lemma 2.** *The cycle  $C$  produced during the second FOR-loop of algorithm  $\mathcal{C}$  does not increase  $d(C^1, C^2, \dots, C^q)$ .*

**Proof of lemma 2.** The proof is done by induction on  $q$ .

#### 4.1.1.1 $q = 1$

The proof of this case is an immediate application of lemma 1 with  $q = 1$ .

#### 4.1.1.2 $q = k$

Suppose that during the  $k$  first executions of the FOR-loop, the number of 2-edges is at most  $k$ .

#### 4.1.1.3 $q = k + 1$

Suppose now that there exists at least one 2-edge in  $C$  (note also that  $C^{s+1}$ , since it has been not processed yet, always contains the 2-edge produced by the execution of the first FOR-loop). Since the patching with  $C^{s+1}$  is done by algorithm  $\mathcal{C}$  using two 2-edges, there is no additional 2-edge created. On the other hand, if no 2-edge exists in  $C$ , the patching of  $C$  with  $C^{s+1}$  will produce at most  $2 \leq k + 1$  new 2-edges and this concludes induction and the proof of lemma 2. ■

Lemmata 1 and 2 induce

$$d(C) \leq c + q \tag{7}$$

### 4.1.2 Construction and evaluation of $D$

The following procedure is used to construct  $D$ .

```

BEGIN /D/
  D ← D1;
  FOR t ← 1 TO r - 1 DO
    replacing as many 2-edges as possible D ← CYCLE_PATCH(D, Dt+1);
  OD
  OUTPUT D;
END. /D/

```

Exactly analogous arguments to the ones of the proof of lemma 2 applied to algorithm  $D$  and thanks to fact 8 induce

$$d(D) \leq d + r \tag{8}$$

Also, let us note that any patching of cycles  $D_i$  between them will create an additional cost of  $r$ .

### 4.1.3 Construction and evaluation of $\tilde{T}$

```

BEGIN / $\tilde{T}$ /
    replacing as many 2-edges as possible  $\tilde{T} \leftarrow \text{CYCLE\_PATCH}(C,D)$ ;
    OUTPUT  $\tilde{T}$ ;
END. / $\tilde{T}$ /

```

With the same arguments as in lemma 2, the following holds for  $|\tilde{T}|$ :

$$\begin{cases} d(\tilde{T}) \leq c + d + q + r & 2q + r \geq 2 \\ d(\tilde{T}) = d & (q, r) = (0, 1) \end{cases} \quad (9)$$

### 4.1.4 Overall specification of the min\_TSP12-algorithm, construction and evaluation of $T$

Once  $\tilde{T}$  constructed, call of  $\text{CYCLE\_PATCH}(\tilde{T}, C_{p+1})$ , changing as many as 2-edges (at most 2) as possible, constructs the final TSP-solution  $T(K_n)$  and the whole min\_TSP12-PTAA is the following. The 2-matching  $\hat{M}$  produced in the first line of the algorithm below is supposed to be optimal and without cycles on less than, or equal to, four edges.

```

BEGIN /TSP12/
    call the algorithm of [14] to produce  $\hat{M}$ ;
     $\hat{M} = (C_1, \dots, C_{p+1}) \leftarrow 2\_MIN(\hat{M})$ ;
     $M^* \leftarrow \text{PREPROCESS}(\hat{M}) \cup_{t=1}^r \{D_t\} \cup C_{p+1}$ ;
     $C \leftarrow C(M^*)$ ;
     $D \leftarrow D(M^*)$ ;
     $\tilde{T} \leftarrow \text{CYCLE\_PATCH}(C,D)$ ;
     $\text{OUTPUT}(K_n) \leftarrow \text{CYCLE\_PATCH}(\tilde{T}, C_{p+1})$ ;
END. /TSP12/

```

It is easy to see that, since all the algorithms called are polynomial, TSP12 works in polynomial time.

If  $(q, r) = (0, 1)$  (in this case, by fact 4,  $C_{p+1} \neq \emptyset$ ), then patching of  $D_1$  with  $C_{p+1}$  constructs a tour with  $d(T(K_n)) = d(D_1) + d(C_{p+1}) + 1 = d(M^*) + 1 = d(M^*) + q + r$ .

Suppose  $2q + r \geq 2$ . Then, by expression (9),  $d(\tilde{T}) \leq c + d + q + r$ . If  $d(\tilde{T}) < c + d + q + r$ , i.e.,  $d(\tilde{T}) \leq c + d + q + r - 1$ , even if  $\tilde{T}$  does not contain any 2-edge, patching of  $\tilde{T}$  with  $C_{p+1}$  will create only one additional 2-edge so, finally,  $d(T(K_n)) \leq d(\tilde{T}) + d(C_{p+1}) \leq c + d + q + r + |C_{p+1}| + |E_2|$  and, by expression (6),  $d(T(K_n)) \leq d(M^*) + q + r$ . If  $d(\tilde{T}) = c + d + q + r$ , we simply exchange two 2-edges and the same expression for  $d(T(K_n))$  always holds.

The discussion above leads to the following concluding expression for the quantity  $d(T(K_n))$ :

$$d(T(K_n)) = \lambda_{\text{TSP12}}(K_n) \leq v(M^*) + q + r \quad (10)$$

## 4.2 A bound for $\omega(K_n)$

In what follows, we will exhibit a TSP12-solution, the objective value of which will provide us with a lower bound for the value  $\omega(K_n)$  of the worst TSP12-solution on  $K_n$ . For this we define a set  $W$  of disjoint elementary paths (d.e.p.), *any one of them containing only 2-edges*. Obviously, if  $W = \{w_1, \dots, w_{|W|}\}$ , one, by properly linking  $w_i$ 's, can easily construct a tour  $T'$  verifying  $d(T') \geq n + \sum_{w_i \in W} |w_i|$  which is a lower bound for  $\omega(K_n)$ .

### 4.2.1 Disjoint elementary paths on $V(C)$

Recall that, for  $q \neq 0$ ,  $d(i^s, I^s) = 1$ ; hence, by fact 1,  $d(a^s, A^s) = d(a^s, B^s) = d(b^s, A^s) = d(b^s, B^s) = 2$ ,  $s = 1, \dots, q$ . Always by fact 1, either  $d(i^s, B^s) = 2$ , or  $d(I^s, b^s) = 2$ . Without loss of generality, we suppose all over the rest of the proof of theorem 2 that  $d(i^s, B^s) = 2$ .

Consequently, for  $s = 1, \dots, q$ , set  $W_{C^s} = \{b^s A^s, A^s a^s, a^s B^s, B^s i^s\}$  and the set of d.e.p. on the vertices of  $C$  is  $W_C = \cup_{s=1}^q W_{C^s}$  with

$$|W_C| = 4q \quad (11)$$

### 4.2.2 Disjoint elementary paths on $V(D)$

If  $r > 1$ , we choose, for  $t = 1, \dots, r$ , a sequence  $\{w_t, x_t, y_t, z_t\} \in V(D_t)$ . Then, the set of d.e.p. and its cardinality on  $V(D)$  is

$$W_D = \{w_1, w_2, \dots, w_t, w_{t+1}, \dots, w_r, x_1, \dots, x_t, \dots, x_r, y_1, \dots, y_r, z_1, \dots, z_r\} \quad (12)$$

$$|W_D| = 4(r-1) + 3 = 4r - 1 \quad (13)$$

If  $r \leq 1$ , then we set  $W_D = \emptyset$ .

### 4.2.3 Disjoint elementary paths on $V(\hat{T})$

#### 4.2.3.1 $q > 0$

Suppose first  $r \neq 1$ . If  $r = 0$ , then  $W_{\hat{T}} = W_C$ . Suppose now  $r > 1$ . Then, by fact 1, there exists vertex  $v_1 \in V(D_1)$  such that either  $d(v_1, B^1) = 2$ , or  $d(v_1, I^1) = 2$ . Let  $e$  be this 2-edge. Without loss of generality, we can suppose  $v_1 = w_1$  (see the paragraph just above). Then the set of d.e.p. on  $V(\hat{T})$  is  $W_{\hat{T}} = W_C \cup W_D \cup \{e\}$  with (see expressions (11) and (13))

$$|W_{\hat{T}}| = 4q + 4r - 1 + 1 = 4(q + r) \quad (14)$$

Suppose  $r = 1$ . Consider cycles  $C_1^1$ ,  $C_2^1$  and  $D_1$  and denote by  $a^{11}$  (resp.,  $A^{11}$ ) the vertex, distinct from  $i^1$  (resp.,  $I^1$ ) in  $C_1^1$  (resp.,  $C_2^1$ ) adjacent to  $a^1$  (resp.,  $A^1$ ). If for any  $u \in V(D_1)$  and for any  $v \in \{a^1, b^1, A^1, B^1\}$ ,  $d(u, v) = 2$ , then let  $w, x, y, z$  be four vertices of  $V(D_1)$  and set  $W_1 = \{wa^1, a^1x, xb^1, b^1y, yA^1, A^1z, zB^1, B^1i^1\}$ .

Suppose now that there exist  $x \in v(D_1)$  and  $v \in \{a^1, b^1, A^1, B^1\}$  with  $d(x, v) = 1$ ; assume  $v = a^1$  (so,  $d(xa^1) = 1$ ). Let  $w, y, z$  be three vertices in  $V(D_1)$  such that  $w, x, y, z$  are subsequent in  $D_1$ . Then, by fact 1,  $d(w, a^{11}) = d(w, i^1) = d(y, a^{11}) = d(y, i^1) = 2$ .

If  $d(y, I^1) = d(x, A^1) = 2$ , then  $W_1 = \{I^1y^1, ya^{11}, a^{11}w, wi^1, a^1B^1, B^1b^1, b^1A^1, A^1x\}$ . If not, we can suppose (up to renaming of cycles  $C_1^1$ ,  $C_2^1$  and  $D_1$  in the discussion that follows)  $d(y, I^1) = 1$ . Then, by fact 1 one of the edges  $i^1x$  and  $a^1y$  is a 2-edge; let us denote it by  $e$ . Set  $f = a^{11}y$  if  $e = i^1x$ , or  $f = i^1w$  if  $e = a^1y$ . Then,  $W_1 = \{a^{11}w, i^1y, a^1A^1, A^1b^1, b^1B^1, B^1z\} \cup \{e, f\}$ . Figure 2 illustrates this case. In all the above cases set  $W_{\hat{T}} = (W_C \setminus W_{C_1}) \cup W_1$  is a set of d.e.p (remark that the hypothesis  $d(i^s, B^s) = 2$  does not intervene in the specification of the set  $W_{\hat{T}}$ ) of cardinality

$$|W_{\hat{T}}| = 4q + 4 + 8 = 4(q + r) \quad (15)$$

From expressions (14) and (15) we conclude for the case  $q > 0$ :

$$|W_{\hat{T}}| = 4(q + r) \quad (16)$$




Figure 2. Construction of  $W_1$ .

#### 4.2.3.2 $q = 0$

Consider first  $|E2| = |C_{p+1}|$ . Remark that  $r \geq 1$  (if not  $T(K_n) = C_{p+1}$  is a minimum-distance Hamiltonian tour); note also that  $C_{p+1}$  can eventually be empty. From facts 2, 3 and 8, for any cycle  $D_t$ ,  $t = 1, \dots, r$  the only 1-edges (other than the ones of  $D_t$ ) incident to vertices of  $V(D_t)$  are pairs of  $V(D_t) \times V(D_t)$  not included in  $D_t$ . However, in any feasible Hamiltonian tour, one cannot use more than  $\sum_{t=1}^{t=r} (|D_t| - 1) = d - r$  of them and, consequently, no less than  $n - (d - r) = |C_{p+1}| + r$  2-edges. Hence,  $\beta(K_n) \geq n + |C_{p+1}| + r = d(M^*) + r = |T(K_n)|$  and the solution computed by algorithm TSP12 is optimal. For case  $|E2| < |C_{p+1}|$ , we set  $W_{\hat{T}} = W_D$  if  $r > 1$ , and  $W_{\hat{T}} = \emptyset$ , if  $r = 1$ .

#### 4.2.4 Disjoint elementary paths on $V(K_n)$

##### 4.2.4.1 $q > 0, |E2| < |C_{p+1}|$

Consider set  $W = W_{\hat{T}} \cup E2$ . Using expression (16), we get  $|W| = 4(q + r) + |E2|$ .

##### 4.2.4.2 $q > 0, |E2| = |C_{p+1}|$

Let  $uv \in C_{p+1}$  and  $u'$  be a vertex with  $|\Gamma_{W_{\hat{T}}}(u')| = 1$ , where by  $\Gamma_{W_{\hat{T}}}(u')$  we denote the set of neighbors of  $u'$  belonging also to  $V(W_{\hat{T}})$ . Remark that such a vertex  $u'$  exists because  $W_{\hat{T}}$  is a simple set of paths. Fact 2 ensures  $d(u, u') = 2$ . We then set  $W = W_{\hat{T}} \cup (C_{p+1} \setminus \{uv\}) \cup \{uu'\}$  with (see expression (16))  $|W| = 4(q + r) + |E2|$ .



#### 4.2.4.3 $q = 0, |E2| < |C_{p+1}|$

Let us first suppose  $r = 1$ . Then, let  $H = \{e_1, e_2, e_3, e_4\}$  be an elementary path on four edges in  $C_{p+1}$  with endpoints  $u$  and  $v$  and such that  $d(e_1) = 1$  and  $d(e_2) = 2$ ; let  $M = \{w, x, y, z\}$  be a sequence of four successive vertices in  $V(D_1)$  and set  $H2 = \{e \in H : d(e) = 2\}$ . Then, using facts 1 and 2, we can construct (see figure 3), between paths  $H$  and  $M$ , a path  $P$  containing at least  $4 + |H2|$  2-edges where  $|\Gamma_P(v)| \leq 1$ . We set  $W = P \cup (E2 \setminus H2)$  that constitutes a d.e.p with  $|W| = (4 + |H2|) + (|E2| - |H2|) = |E2| + 4$ .




Figure 3. Construction of  $W$  supposing  $vz = \operatorname{argmax}\{d(v', y), d(v, z)\}$ .

Let us now suppose  $r > 1$  and let  $uv$  be an 1-edge of  $C_{p+1}$ . Moreover, from the previous paragraph, for the case we deal with,  $W_{\hat{T}} = W_D$ , where  $W_D$  is given by expression (12). By fact 1 we have that either  $x_1u$ , or  $y_1v$  is a 2-edge; let us suppose  $d(x_1, u) = 2$ . Then, the set  $W = W_{\hat{T}} \cup E2 \cup \{x_1u\}$  forms a d.e.p. composed of  $|W| = (4r - 1) + |E2| + 1 = 4r + |E2| = 4(q + r) + |E2|$  2-edges.

Consequently, dealing with  $W$ , we always have  $|W| \geq 4(q + r) + |E2|$ . One can obtain a tour  $T_w(K_n)$  by properly linking d.e.ps by edges (at worst by 1-edges) in order that they form a Hamiltonian cycle on  $K_n$ . The so obtained  $T_w(K_n)$  has objective value  $d(T_w(K_n)) \geq n + 4(q + r) + |E2|$ ; so, using expression (6)

$$\omega(K_n) \geq d(T_w(K_n)) \geq n + 4(q + r) + |E2| = d(M^*) + 4(q + r). \quad (17)$$

### 4.3 The differential approximation ratio of TSP12

We have already seen that if  $q = 0$  and  $|E2| = |C_{p+1}|$ , then  $\delta(\min\_TSP12) = 1$ . So, for  $q > 0$  or  $q = 0$  and  $|E2| < |C_{p+1}|$  expressions (10), (17) and the fact that  $\beta(K_n) \geq d(M^*)$ , we get

$$\delta_{TSP12}(K_n) = \frac{\omega(K_n) - \lambda_{TSP12}(K_n)}{\omega(K_n) - \beta(K_n)} \geq \frac{d(M^*) + 4(q+r) - (d(M^*) + (q+r))}{d(M^*) + 4(q+r) - d(M^*)} = \frac{3(q+r)}{4(q+r)} = \frac{3}{4}.$$

### 4.4 Ratio 3/4 is tight for TSP12




Figure 4. Tightness of the TSP12 approximation ratio.

Consider two cliques and number their vertices by  $\{1, \dots, 4\}$  and by  $\{5, 6, \dots, n+8\}$ , respectively. Edges of both cliques have all distance 1. Cross-edges  $ij$ ,  $i = 1, 3$ ,  $j = 5, \dots, n+8$ , are all of distance 2, while every other cross-edge is of distance 1.

Unraveling of TSP12 will produce:

$$\begin{aligned} T &= \{1, 2, 3, 4, 5, 6, \dots, n+7, n+8, 1\} && \text{cycle-pathing on edges } (1, 4) \text{ and } (5, n+8) \\ T_w &= \{1, 5, 2, 6, 3, 7, 4, 8, 9, \dots, n+7, n+8, 1\} && \text{using 2-edges } (1, 5), (6, 3), (3, 7) \text{ and } (n+8, 1) \\ T^* &= \{1, 2, n+8, n+7, \dots, 5, 4, 3, 1\} && \text{using 1-edges } (4, 5), (2, n+8) \end{aligned}$$

i.e.,  $\lambda(K_{n+8}) = n+9$ ,  $\beta(K_{n+8}) = n+8$  and  $\omega(K_{n+8}) = n+12$  (in figure 4,  $T^*$  and  $T_w$  are shown for  $n=2$ ;  $T = \{1, \dots, 10, 1\}$ ). Consequently,  $\delta_{TSP12}(K_{n+8}) = 3/4$  and this completes the proof of theorem 2. ■

Let us note that the differential approximation ratio of the 7/6-algorithm of [18], when running on  $K_{n+8}$ , is also 3/4. The authors of [18] bring also to the fore a family of worst-case instances for their algorithm: one has  $k$  cycles of length four arranged around a cycle of length  $2k$ . We have performed a limited comparative study between their algorithm and the our one, for  $k = 3, 4, 5, 6$  (on 24 graphs). The average differential and standard approximation ratios for the two algorithms are presented in table 1.

## 5 Further results for minimum traveling salesman

### 5.1 Bridges between differential and standard approximation

Let us consider the following approximation-preserving reduction proposed in [16], strongly inspired by the  $A$ -reduction of [17] between pairs  $(\Pi, R)$ , where  $\Pi$  is an **NPO** problem and  $R$  an

	$k$	TSP12	The algorithm of [18]
Differential ratio	3	0,931100364	0,846702091
	4	0,9000002	0,833333
	5	0,920289696	0,833333
	6	0,9222222	0,833333
Standard ratio	3	0,923350955	0,87013
	4	0,9094018	0,857143
	5	0,92646313	0,857143
	6	0,928178	0,857143

Table 1: A limited comparison between TSP12 and the algorithm of [18] on some worst-case instances of the latter.

approximation measure. In what follows, we denote by  $R[\Pi](I, S)$  the value of the approximation measure  $R$  relative to a solution  $S$  of an instance  $I$  of  $\Pi$ . We suppose that  $R$  has values in  $[0, 1]$  (for the standard approximation, we inverse the approximation ratio in the case of minimization problems).

**Definition 3.** A  $G$ -reduction of the pair  $(\Pi_1, R_1)$  to  $(\Pi_2, R_2)$ , denoted by  $(\Pi_1, R_1) \leq^G (\Pi_2, R_2)$ , is a triplet  $(\alpha, g, c)$  such that:

- $\alpha: \mathcal{I}_1 \rightarrow \mathcal{I}_2$  polynomially transforms instances of  $\Pi_1$  into instances of  $\Pi_2$ ;
- $g: \mathcal{S}(\alpha(I)) \rightarrow \mathcal{S}(I)$  polynomially transforms solutions for  $\Pi_2$  into solutions for  $\Pi_1$ ;
- $c: [0, 1] \rightarrow [0, 1]$  ( $c^{-1}(0) = \{0\}$ ) is such that,  $\forall \epsilon \in [0, 1], \forall I \in \mathcal{I}_1, \forall S \in \mathcal{S}(\alpha(I))$ ,

$$R_2[\Pi_2](\alpha(I), S) \geq \epsilon \implies R_1[\Pi_1](I, g(S)) \geq c(\epsilon). \quad \blacksquare$$

The following easy lemma holds.

**Lemma 3.** Consider an **NPO** problem  $\Pi = (\mathcal{I}, \mathcal{S}, v_I, \text{opt})$ . If  $\exists t > 0$  such that,  $\forall I \in \mathcal{I}$ ,  $|\omega(I) - \beta(I)| \leq t \min\{\omega(I), \beta(I)\}$ , then  $(\Pi, \rho) \leq^G (\Pi, \delta)$  with

$$c_t(\epsilon) = \begin{cases} \frac{t\epsilon+1}{t+1} & \text{opt} = \max \\ \frac{1}{t+1-t\epsilon} & \text{opt} = \min \end{cases}$$

Remark that for  $\min\_TSPab$  we have

$$\omega(K_n) - \beta(K_n) \leq bn - an \leq (b-a)n \leq \frac{b-a}{a}\beta(K_n)$$

and by application of lemma 3 the following theorem holds.

**Theorem 3.**

$$\begin{aligned} (\min\_TSPab, \rho) &\leq^G (\min\_TSPab, \delta) \text{ with } c(\epsilon) = \frac{a}{b - (b-a)\epsilon} \\ (\min\_TSP12, \rho) &\leq^G (\min\_TSP12, \delta) \text{ with } c(\epsilon) = \frac{1}{2-\epsilon}. \end{aligned}$$

Theorem 3 implies  $1/\rho_{\text{TSP12}} \geq 4/5$ ; in other words,  $\rho_{\text{TSP12}} \leq 5/4$ . This ratio is better than the one of [5] for this particular case, but with no operational impact since it is dominated by the result of [18].

Recall that  $\min\_TSP12$  and  $\min\_TSPab$  are equi-approximable in the differential approximation framework. Consequently, using theorem 3 with  $\delta = 3/4$ , the following corollary holds.

**Corollary 3.**  *$\min\_TSPab$  is approximable within*

$$\rho \leq \frac{3}{4} + \frac{1}{4} \frac{b}{a}$$

*in the standard framework. This ratio tends to  $\infty$  with  $b$ .*

Let us now denote by  $A_{\max}$  and  $A_{\min}$  a maximum and a minimum spanning trees of  $K_n$ , respectively, and by  $c(A_{\max})$  and  $c(A_{\min})$  their respective costs. Then, the following proposition holds.

**Proposition 4.** *If  $c(A_{\max})/c(A_{\min}) \leq \nu$ ,  $\nu > 1$ , then  $(\min\_TSP, \rho) \leq^G (\min\_TSP, \delta)$  with  $c(\epsilon) = 1/(\nu(1 - \epsilon) + \epsilon)$ .*

**Proof.** Let  $T_w(K_n)$  and  $T^*(K_n)$  be a worst-value tour and an optimal tour of  $K_n$ , respectively. Set  $d_w = \min_{v_i, v_j \in T_w(K_n)} \{d(i, j)\}$  and  $d_\beta = \max_{v_i, v_j \in T^*(K_n)} \{d(i, j)\}$ . Since  $T_w(K_n) \setminus \{\text{argmin}_{v_i, v_j \in T_w(K_n)} \{d(i, j)\}\}$  and  $T^*(K_n) \setminus \{\text{argmax}_{v_i, v_j \in T^*(K_n)} \{d(i, j)\}\}$  are obviously spanning trees of  $K_n$ :  $c(A_{\max}) \geq \omega(K_n) - d_w$ ,  $c(A_{\min}) \leq \beta(K_n) - d_\beta$ . Remark also that  $d_w \leq \omega(K_n)/n$  and  $d_\beta \geq \beta(K_n)/n$ . So,  $c(A_{\max}) \geq \omega(K_n)(1 - 1/n)$  and  $c(A_{\min}) \leq \beta(K_n)(1 - 1/n)$ . Consequently,

$$\frac{\omega(K_n)}{\beta(K_n)} \leq \frac{c(A_{\max})(1 - \frac{1}{n})}{c(A_{\min})(1 - \frac{1}{n})} \leq \frac{c(A_{\max})}{c(A_{\min})} \leq \nu.$$

Hence,  $\omega(K_n) - \beta(K_n) \leq (\nu - 1)\beta(K_n)$ , and using lemma 3 for  $t = (\nu - 1)$  we get  $c(\epsilon) = (\nu(1 - \epsilon) + \epsilon)^{-1}$ . ■

## 5.2 An inapproximability result

We first note that one can prove very easily (with arguments similar to the ones of theorem 6.13 in [13]) that  $\min\_TSP$  cannot be solved by a differential FPTAS unless  $\mathbf{P} = \mathbf{NP}$ . We now restrict ourselves to  $\min\_TSP12$  and revisit theorem 3. It is easy to see that it does not only establish links between the approximabilities of  $\min\_TSP12$  in standard and differential frameworks, but it also establishes limits on its approximability in the two frameworks. Plainly, since approximation of  $\min\_TSP12$  within  $\delta = 1 - \epsilon$  implies its approximation within  $\rho = 2 - (1 - \epsilon) = 1 + \epsilon$ ,  $0 \leq \epsilon \leq 1$ , if there exists an  $\epsilon_0$  such that, under a very likely complexity theory hypothesis,  $\min\_TSP12$  is inapproximable within  $\rho_0 \leq 1 + \epsilon_0$ , then it is inapproximable within  $\delta_0 \geq 1 - \epsilon_0$ . In other words, the hardness thresholds for standard and differential frameworks are identical.

**Theorem 4.** *If under a complexity theory hypothesis  $\min\_TSP12$  is inapproximable within  $1 + \epsilon_0$ , then, under the same hypothesis,  $\min\_TSP12$  is differentially inapproximable within  $1 - \epsilon_0$ .*

Recall the negative result of [4]:  $\forall \epsilon > 0$ , no PTAA can guarantee standard approximation ratio less than, or equal to,  $3477/3476 - \epsilon$  unless  $\mathbf{P} = \mathbf{NP}$ . Using theorem 4,  $\forall \epsilon > 0$ , it is  $\mathbf{NP}$ -hard to approximate  $\min\_TSP12$  with differential ratio better than  $3475/3476 + \epsilon$ . Since  $\min\_TSP12$  is a special case of general  $\min\_TSP$ , the following corollary holds concluding the section.

**Corollary 4.**  *$\min\_TSP$  cannot be approximated within differential ratio greater than, or equal to,  $3475/3476 + \epsilon$ , for every positive  $\epsilon$ , unless  $\mathbf{P} = \mathbf{NP}$ .*

Finally, let us note that the inapproximability result of [11] for dense graphs holds also in the differential approximation framework with the same hardness threshold.

## 6 Differential approximation of maximum traveling salesman

We have also mentioned that in the opposite of  $\min\_TSP$ ,  $\max\_TSP$  (certainly less popular than its cousin), although it is **APX**-hard ([20, 18]), can be solved by a PTAA achieving standard approximation ratio  $\rho = 5/7$  (this ratio is somewhat worst –  $38/63$  – when the input-graph is directed).

The purpose of this section is to show that, in the differential approximation framework, the two cousins are equi-approximable establishing so a kind of natural symmetry between the two problems at hand.

**Theorem 5.**  *$\max\_TSP$  is equi-approximable with  $\min\_TSP$ ; consequently it is in **D-APX**.*

**Proof.** Observe first that, given a graph  $K_n$ , there exists a very interesting symmetry between  $\min\_TSP$  and  $\max\_TSP$  with respect to worst-case and best objective values:

$$\begin{cases} \beta_{\min}(K_n) = \omega_{\max}(K_n) \\ \beta_{\max}(K_n) = \omega_{\min}(K_n) \end{cases} \quad (18)$$

Expression (18) confirms what we said in the introduction of the paper that the worst value of a problem can be as hard to compute as the optimal one.

Given a complete graph  $K_n$ , let us denote by  $\bar{K}_n$  the complete graph on  $n$  vertices when one replaces distance  $d(i, j)$  by  $\bar{d}(i, j) = M - d(i, j)$ ,  $i, j = 1, \dots, n$ , for  $M = \max_{v_i, v_j \in E} \{d(i, j)\} + \min_{v_i, v_j \in E} \{d(i, j)\}$ . It is easy to see that  $\bar{\bar{K}}_n = K_n$ . Moreover, any TSP-feasible solution for  $K_n$  is TSP-feasible for  $\bar{K}_n$ .

Given a Hamiltonian cycle  $T$ , we use notation  $T_{\min}$  (resp.,  $T_{\max}$ ) in order to indicate that we deal with a solution of  $\min\_TSP$  (resp.,  $\max\_TSP$ ). We then have

$$\begin{aligned} |T_{\min}(K_n)| &= Mn - |T_{\max}(\bar{K}_n)| \\ |T_{\max}(K_n)| &= Mn - |T_{\min}(\bar{K}_n)| \end{aligned}$$

and, more particularly,

$$\omega_{\min}(K_n) = Mn - \beta_{\min}(\bar{K}_n) = Mn - \omega_{\max}(\bar{K}_n) \quad (19)$$

$$\beta_{\min}(K_n) = Mn - \omega_{\min}(\bar{K}_n) = Mn - \beta_{\max}(\bar{K}_n) \quad (20)$$

$$\lambda_{\mathbf{A}}^{\min}(K_n) = Mn - \lambda_{\mathbf{A}}^{\max}(\bar{K}_n) \quad (21)$$

By the discussion above, one can immediately conclude that for every PTAA  $\mathbf{A}$  and for every  $K_n$ ,  $\delta_{\mathbf{A}}^{\min}(K_n) = \delta_{\mathbf{A}}^{\max}(\bar{K}_n)$  (where, once again, indices  $\min$  and  $\max$  are used to denote  $\min\_TSP$  and  $\max\_TSP$ , respectively). Consequently,  $\delta_{\mathbf{A}}^{\min} = \delta_{\mathbf{A}}^{\max}$ ,  $\forall \mathbf{A}$ . Since  $\delta_{2\_OPT}^{\min} \geq 1/2$ , the same holds for  $\delta_{2\_OPT}^{\max}$  and this completes the proof of the theorem. ■

For  $d(i, \bar{j}) \in \{a, b\}$ ,  $\max_{v_i, v_j \in E} \{d(i, j)\} + \min_{v_i, v_j \in E} \{d(i, j)\} - d(i, j) \in \{a, b\} \forall v_i, v_j \in E$ ; so, the proof of theorem 5 establishes also equi-approximability between  $\min\_TSPab$  and  $\max\_TSPab$  and the following theorem summarizes differential approximation results for  $\max\_TSP$ .

**Theorem 6.**

- $\max\_TSP$  is approximable within differential approximation ratio  $1/2$ ;
- $\max\_TSP12$  and  $\max\_TSPab$  are approximable within differential approximation ratio  $3/4$ ;
- for every  $\epsilon > 0$ ,  $\max\_TSP$  cannot be approximated within differential ratio greater than, or equal to,  $5379/5380 + \epsilon$ , unless **P=NP**.

## 7 An improvement of the standard ratio for the maximum traveling salesman with distances 1 and 2

Application of lemma 3 in the case of  $\max\_TSPab$  with  $t = (b - a)/a$  gets

$$\rho = c_{\frac{b-a}{a}}(\delta) = \frac{b-a}{b}\delta + \frac{a}{b}$$

and for  $\delta = 3/4$  we have

$$\rho = c_{\frac{b-a}{a}}\left(\frac{3}{4}\right) = \frac{3}{4} + \frac{1}{4}\frac{a}{b} \quad (22)$$

The above ratio is always bounded below by  $3/4$ . Here we see another impact of the asymmetry between minimization and maximization versions of TSP in the standard approximation framework. Recall that, as we have seen in section 5.1, the standard approximation ratio for  $\min\_TSPab$  tends to  $\infty$  with  $b$  and this obviously holds for every PTAA.

Set now  $a = 1$  and  $b = 2$  and revisit expression (22). Then, the following theorem immediately holds.

**Theorem 7.**  *$\max\_TSP12$  is polynomially approximable within  $\rho^{\max} \geq 7/8$ .*

Such an improved ratio ( $7/8 > 5/7$ ) for  $\max\_TSP12$  seems that it cannot be immediately achieved by the interesting work of [20].

Consider now the following algorithm for  $\max\_TSP$ .

```

BEGIN /MTSPALG/
  construct  $\bar{K}_n$ ;
  call the algorithm of [18] to compute a tour  $T_{\min}(\bar{K}_n)$ ;
  OUTPUT  $T_{\max}(K_n) \leftarrow T_{\min}(\bar{K}_n)$ ;
END.
```

Recall that the algorithm called in the first line of the algorithm just above guarantees  $\rho^{\min} \leq 7/6$ . Then, using expressions (20), (21) and some easy algebra, one gets  $\rho_{\text{MTSPALG}}^{\max}(K_n) \geq 2/3$ .

## 8 Towards stronger differential-inapproximability results for the traveling salesman

Recall that an **NPO** problem  $\Pi$  is called *simple* ([19]) if its restriction  $\Pi_k$  to instances verifying, for every fixed constant  $k \in \mathbb{N}$ ,  $\beta(I) \leq k$  can be solved in polynomial time. Analogously, we will call  $\Pi$  *D-simple* if its restriction  $\Pi_k$  to instances verifying, for every fixed constant  $k \in \mathbb{N}$ ,  $|\omega(I) - \beta(I)| \leq k$  is polynomial. Then the following proposition holds.

**Proposition 5.** *If  $\Pi$  is not D-simple, then there exists  $k_0 \in \mathbb{N}$  such that  $\delta(\Pi) < k_0/(k_0 + 1)$ .*

**Proof.** Suppose  $\Pi$  not *D-simple* and  $\delta(\Pi) \geq k/(k + 1)$ ,  $\forall k \in \mathbb{N}$ . Then,  $\forall I \in \mathcal{I}$ ,

$$\omega(I) - \lambda(I) \geq \omega(I) - \beta(I) - \frac{\omega(I) - \beta(I)}{k + 1} \quad (23)$$

Consider now an instance  $I' \in \mathcal{I}$  such that  $\omega(I') - \beta(I') \leq k$ . Then, since objective function's values are integer (item 3 of definition 1), expression (23) gives  $\omega(I') - \lambda(I') = \omega(I') - \beta(I')$ . Consequently, it suffices to set  $k_0 = \min\{k : \Pi_k \text{ non polynomial}\}$  in order to complete the proof. ■

In other words, problems which are not *D-simple* do not admit differential PTAS (this is the differential-equivalent of the result of [19] for the standard approximation). The proposition above allows achievement of stronger hardness thresholds provided that  $k_0$  is a fixed constant. We conjecture that TSP is not *D-simple* and this for a small  $k_0$ . If this was true, the hardness threshold of corollary 4 could be meaningfully improved.

## References

- [1] S. Arora. Polynomial time approximation schemes for Euclidean TSP and other geometric problems. In *Proc. FOCS'96*, pages 2–11, 1996.
- [2] S. Arora, C. Lund, R. Motwani, M. Sudan, and M. Szegedy. Proof verification and intractability of approximation problems. In *Proc. FOCS'92*, pages 14–23, 1992.
- [3] G. Ausiello, A. D'Atri, and M. Protasi. Structure preserving reductions among convex optimization problems. *J. Comput. System Sci.*, 21:136–153, 1980.
- [4] H. Böckenhauer, J. Hromkovič, R. Klasing, S. Seibert, and W. Unger. An improved lower bound on the approximability of metric TSP and approximation algorithms for the TSP with sharpened triangle inequality. In *Proc. STACS'2000*. To appear.
- [5] N. Christofides. Worst-case analysis of a new heuristic for the traveling salesman problem. Technical Report 388, Grad. School of Industrial Administration, CMU, 1976.
- [6] P. Crescenzi and V. Kann. A compendium of NP optimization problems. Available on `www_address: http://www.nada.kth.se/~viggo/problemlist/compendium.html`, 1995.
- [7] A. Croes. A method for solving traveling-salesman problems. *Oper. Res.*, 5:791–812, 1958.
- [8] M. Demange, P. Grisoni, and V. T. Paschos. Differential approximation algorithms for some combinatorial optimization problems. *Theoret. Comput. Sci.*, 209:107–122, 1998.
- [9] M. Demange and V. T. Paschos. On an approximation measure founded on the links between optimization and polynomial approximation theory. *Theoret. Comput. Sci.*, 158:117–141, 1996.
- [10] L. Engebretsen. An explicit lower bound for TSP with distances one and two. Technical Report TR 98-046, revised, Electronic Colloquium on Computational Complexity, January 1999. Available on `www_address: http://www.eccc.uni-trier.de/eccc/`.
- [11] W. Fernandez de la Vega and M. Karpinski. On approximation hardness of dense TSP and other path problems. *Inform. Process. Lett.*, 70:53–55, 1999.
- [12] M. L. Fisher, G. L. Nemhauser, and L. A. Wolsey. An analysis of approximations for finding a maximum weight Hamiltonian circuit. *Oper. Res.*, 27(4):799–809, 1979.
- [13] M. R. Garey and D. S. Johnson. *Computers and intractability. A guide to the theory of NP-completeness*. W. H. Freeman, San Francisco, 1979.
- [14] D. B. Hartvigsen. *Extensions of matching theory*. PhD thesis, Carnegie-Mellon University, 1984.
- [15] S. Lin and B. W. Kernighan. An effective heuristic algorithm for the traveling salesman problem. *Oper. Res.*, 21:498–516, 1973.
- [16] J. Monnot. Families of critical instances and polynomial approximation. PhD Thesis. LAMSADE, Université Paris-Dauphine, 1998 (in French).
- [17] P. Orponen and H. Mannila. On approximation preserving reductions: complete problems and robust measures. Technical Report C-1987-28, Dept. of Computer Science, University of Helsinki, Finland, 1987.

- [18] C. H. Papadimitriou and M. Yannakakis. The traveling salesman problem with distances one and two. *Math. Oper. Res.*, 18:1–11, 1993.
- [19] A. Paz and S. Moran. Non deterministic polynomial optimization problems and their approximations. *Theoret. Comput. Sci.*, 15:251–277, 1981.
- [20] S. Rao Kosaraju, J. K. Park, and C. Stein. Long tours and short superstrings. In *Proc. FOCS'94*, pages 166–177, 1994.
- [21] S. Sahni and T. Gonzales. P-complete approximation problems. *J. Assoc. Comput. Mach.*, 23:555–565, 1976.
- [22] S. Vishwanathan. An approximation algorithm for the asymmetric travelling salesman problem with distances one and two. *Inform. Process. Lett.*, 44:297–302, 1992.
- [23] E. Zemel. Measuring the quality of approximate solutions to zero-one programming problems. *Math. Oper. Res.*, 6:319–332, 1981.